

BLUE BIRD
DCX Sheep & Goat Feed/Milk Replacer
 (decoquinatate Type C medicated feed)

Sheep: For the prevention of coccidiosis in young sheep caused by *Eimeria ovinoidalis*, *E. parva*, *E. bakuensis*, and *E. crandallis*.

Goats: For the prevention of coccidiosis in young goats caused by *Eimeria christensenii* and *E. ninakohlyakimovae*.

Active Drug Ingredient

Decoquinatate..... 12.9 to 90.8 g/ton*

Guaranteed Analysis

Crude Protein (Min).....	_____ %
NPN ^a (Max).....	_____ %
Crude Fat (Min).....	_____ %
Crude Fiber (Max).....	_____ %
Calcium ^a (Min).....	_____ %
Calcium ^a (Max).....	_____ %
Phosphorus ^a (Min).....	_____ %
Salt ^a (Min).....	_____ %
Salt ^a (Max).....	_____ %
Sodium ^b (Min).....	_____ %
Sodium ^b (Max).....	_____ %
Copper ^{c, d} (Min).....	_____ ppm or %
Copper ^{c, d} (Max).....	_____ ppm or %
Selenium ^{a, d} (Min).....	_____ ppm or %
Vitamin A ^a (Min).....	_____ IU/lb

^a Guarantee required only when nutrient added except when the feed is intended, represented or serves as a principal source of the nutrient.

^b Sodium guarantee required only when total sodium exceeds that furnished by the maximum salt guarantee.

^c Copper minimum and maximum guarantee for sheep required when added or level exceeds 20 ppm.

^d Guarantee shall be stated in ppm when less than 10,000 ppm and in percentage when concentration is 10,000 ppm (1%) or greater.

Ingredients

Ingredients as defined by AAFCO.

Feeding Directions

Feed at a rate of 22.7 mg decoquinatate per 100 lb of body weight (0.5 mg/kg) per day. Feed at least 28 days during period of exposure to coccidiosis or when it is likely to be a hazard.

Warning: Do not feed to sheep or goats producing milk for human consumption.

Caution: Do not use in feeds containing bentonite.

Manufactured by:

Blue Bird Feed Mill

City, State, Zip

NET WEIGHT ON BAG OR BULK

Expiration Date (if applicable) _____

Lot Number (if applicable)) _____

*Label must show actual concentration of decoquinatate, which will vary with body weight and feed consumption to provide 22.7 mg/100 lb body weight per day. The following table provides examples (100% dry matter basis) of feeding rates and amount of drug per ton of feed.

Animal Wt (Lbs.)	Feed Intake (% Body Wt)	Lbs. Feed/Head/Day	Decoquinatate (g/ton)
25	0.5	0.125	90.8
	1.5	0.375	30.3
	3.5	0.875	13.0
50	0.5	0.25	90.8
	1.5	0.75	30.3
	3.5	1.75	13.0
100	0.5	0.50	90.8
	1.5	1.50	30.3
	3.5	3.50	13.0