

**BLUE BIRD**  
**DCX Top-dress Sheep & Goat Feed**  
(decoquinate Type C top-dress medicated feed)

**Sheep:** For the prevention of coccidiosis in young sheep caused by *Eimeria ovinoidalis*, *E. parva*, *E. bakuensis*, and *E. crandallis*.

**Goats:** For the prevention of coccidiosis in young goats caused by *Eimeria christensenii* and *E. ninakohlyakimovae*.

**Active Drug Ingredient**

Decoquinate.....90.9 – 535.7 g/ton\*

**Guaranteed Analysis**

|                                     |       |          |
|-------------------------------------|-------|----------|
| Crude Protein (Min).....            | _____ | %        |
| NPN <sup>a</sup> (Max).....         | _____ | %        |
| Crude Fat (Min).....                | _____ | %        |
| Crude Fiber (Max).....              | _____ | %        |
| Calcium <sup>a</sup> (Min).....     | _____ | %        |
| Calcium <sup>a</sup> (Max).....     | _____ | %        |
| Phosphorus <sup>a</sup> (Min).....  | _____ | %        |
| Salt <sup>a</sup> (Min).....        | _____ | %        |
| Salt <sup>a</sup> (Max).....        | _____ | %        |
| Sodium <sup>b</sup> (Min).....      | _____ | %        |
| Sodium <sup>b</sup> (Max).....      | _____ | %        |
| Copper <sup>c, d</sup> (Min).....   | _____ | ppm or % |
| Copper <sup>c, d</sup> (Max).....   | _____ | ppm or % |
| Selenium <sup>a, d</sup> (Min)..... | _____ | ppm or % |
| Vitamin A <sup>a</sup> (Min).....   | _____ | IU/lb    |

<sup>a</sup> Guarantee required only when nutrient added except when the feed is intended, represented or serves as a principal source of the nutrient.

<sup>b</sup> Sodium guarantee required only when total sodium exceeds that furnished by the maximum salt guarantee.

<sup>c</sup> Copper minimum and maximum guarantee for sheep required when added or level exceeds 20 ppm.

<sup>d</sup> Guarantee shall be stated in ppm when less than 10,000 ppm and in percentage when concentration is 10,000 ppm (1%) or greater.

**Ingredients**

Ingredients as defined by AAFCO.

**Feeding Directions**

Administer as a top dress supplement or mix into the daily ration to provide 22.7 mg per 100 pounds of body weight (0.5 mg/kg) per day. \_\_\_\_\_\*\* pounds of this supplement will medicate 100 lbs. of body weight. Feed at least 28 days during period of exposure to coccidiosis or when it is likely to be a hazard.

**Warning:** Do not feed to sheep and goats producing milk for human consumption.

**Caution:** Do not use in feeds containing bentonite.

**Manufactured by:**

Blue Bird Feed Mill

City, State, Zip

**NET WEIGHT ON BAG OR BULK**

Expiration Date (if applicable)\_\_\_\_\_

Lot Number (if applicable) )\_\_\_\_\_

\*Label must show actual concentration of decoquinate, which will vary with body weight and feed consumption to provide 22.7 mg/100 lb body weight per day.

\*\*The following table provides examples of supplementation rates and amount of drug per ton of feed. Third column represents number of pounds of supplement that will medicate 100 lbs of body weight.

| Drug level in supplement |       | ** Lbs. supplement/<br>100 lbs. body weight |
|--------------------------|-------|---|
| mg/lb                    | g/ton |   |
| 45.45                    | 90.9  | 0.5   |
| 56.75                    | 113.5 | 0.4   |
| 113.5                    | 227   | 0.2   |
| 227                      | 454   | 0.1   |
| 267                      | 534   | 0.085                                       |