

**Label 1: Broiler Chickens Type C Medicated Feed Coccidiostat**

**Indications for Use**

For the prevention of coccidiosis caused by *E. tenella* at the level of 72.6 – 113.5 g/ton (0.008 - 0.0125%)

**Active Drug Ingredient**

Amprolium.....0.008-0.0125%

**Guaranteed Analysis**

Crude Protein, not less than . . . . .xx.x%  
Crude Fat, not less than . . . . .xx.x%  
Crude Fiber, not more than . . . . .xx.x%  
Lysine, not less than . . . . .xx.x%  
Methionine, not less than . . . . .xx.x%  
Calcium, min and max . . . . .xx.x%  
Phosphorus, not less than . . . . .xx.x%  
Salt, min and max (if added) . . . . .xx.x%  
Sodium, min and max (when total Sodium exceed the  
furnished by the maximum salt guarantee) . . . . .xx.x%

**Ingredients**

Ingredients as defined by AAFCO

**Feeding Directions**

**Prevention:**

For the prevention of coccidiosis caused by *E. tenella* only, feed 0.008 – 0.0125% amprolium Type C medicated feed.

Increasing levels of Amprolium may not prevent coccidiosis caused by strains of *Eimeria* commonly found to be resistant to amprolium.

**Caution**

Use as the sole source of amprolium.

Do not change the litter while giving this feed unless absolutely necessary.

If losses exceed 0.5% in a 2-day period, obtain an accurate diagnosis, and follow the instructions of your veterinarian or poultry pathologist.

Losses may result from intercurrent disease or other conditions affecting drug intake which can contribute to the virulence of coccidiosis under field conditions.

Do not use in feeds containing bentonite.

**Warning**

**Keep this and all drugs out of the reach of children.**

**Manufactured By**

Blue Bird Feed Mill  
Robin, Indiana 00000

Net Weight \_\_\_ lbs (\_\_\_ kg)  
Bag or Bulk