

Label 2: Broiler Chickens Type C Medicated Feed Coccidiostat

Indications for Use

For the prevention of coccidiosis where immunity is not desired at the level of 113.5-227 g/ton (0.0125% - 0.025%)

Active Drug Ingredient

Amprolium.....0.0125-0.025%

Guaranteed Analysis

Crude Protein, not less thanxx.x%
Crude Fat, not less thanxx.x%
Crude Fiber, not more thanxx.x%
Lysine, not less thanxx.x%
Methionine, not less thanxx.x%
Calcium, min and maxxx.x%
Phosphorus, not less thanxx.x%
Salt, min and max (if added)xx.x%
Sodium, min and max (when total Sodium exceed the
furnished by the maximum salt guarantee)xx.x%

Ingredients

Ingredients as defined by AAFCO

Feeding Directions

Prevention:

For most field conditions under modern management practices and where immunity is not desired, feed 0.0125% amprolium. Where severe coccidiosis conditions exist, and immunity is not desired, use 0.025% amprolium Type C medicated feed.

Increasing levels of Amprolium may not prevent coccidiosis caused by strains of *Eimeria* commonly found to be resistant to amprolium.

Caution

Use as the sole source of amprolium.

Do not change the litter while giving this feed unless absolutely necessary.

If losses exceed 0.5% in a 2-day period, obtain an accurate diagnosis, and follow the instructions of your veterinarian or poultry pathologist.

Losses may result from intercurrent disease or other conditions affecting drug intake which can contribute to the virulence of coccidiosis under field conditions.

Do not use in feeds containing bentonite.

Warning

Keep this and all drugs out of the reach of children.

Manufactured By

Blue Bird Feed Mill
Robin, Indiana 00000

Net Weight ___ lbs (___ kg)
Bag or Bulk