

Label 2: Replacement Chickens Type C Medicated Feed Coccidiostat	
Indications for Use	
For the prevention of coccidiosis where immunity to coccidiosis is not desired.	
Active Drug Ingredient	
Amprolium.....	0.0125-0.025%
Guaranteed Analysis	
Crude Protein, not less thanxx.x%
Crude Fat, not less than	xx.x%
Crude Fiber, not more thanxx.x%
Lysine, not less than	xx.x%
Methionine, not less than	xx.x%
Calcium, min and max	xx.x%
Phosphorus, not less thanxx.x%
Salt, min and max (if added)xx.x%
Sodium, min and max (when total Sodium exceed the furnished by the maximum salt guarantee)	xx.x%
Ingredients	
Ingredients as defined by AAFCO	
Feeding Directions	
Feed continuously from day old until onset of production.	
Caution	
Use as the sole source of amprolium.	
Do not change the litter while giving this feed unless absolutely necessary.	
If losses exceed 0.5% in a 2-day period, obtain an accurate diagnosis, and follow the instructions of your veterinarian or poultry pathologist.	
Losses may result from intercurrent disease or other conditions affecting drug intake which can contribute to the virulence of coccidiosis under field conditions.	
In replacement flocks the grower must expect that excessive exposure to one or more species may overwhelm the drug in some flocks and prompt treatment will be required.	
Do not use in feeds containing bentonite.	
Warning	
Keep this and all drugs out of the reach of children.	
Manufactured By	
Blue Bird Feed Mill Robin, Indiana 00000	
Net Weight ___ lbs (___ kg) Bag or Bulk	