

Turkeys Type C Medicated Feed Coccidiostat

Indications for Use

To be used in prevention of coccidiosis.

Active Drug Ingredient

Amprolium.....0.0125-0.025%

Guaranteed Analysis

Crude Protein, not less thanxx.x%
Crude Fat, not less thanxx.x%
Crude Fiber, not more thanxx.x%
Lysine, not less thanxx.x%
Methionine, not less thanxx.x%
Calcium, min and maxxx.x%
Phosphorus, not less thanxx.x%
Salt, min and max (if added)xx.x%
Sodium, min and max (when total Sodium exceed the
furnished by the maximum salt guarantee)xx.x%

Ingredients

Ingredients as defined by AAFCO

Feeding Directions

Feed continuously.

Caution

Use as the sole source of amprolium.

Do not change the litter while giving this feed unless absolutely necessary.

If losses exceed 0.5% in a 2-day period, obtain an accurate diagnosis, and follow the instructions of your veterinarian or poultry pathologist.

Losses may result from intercurrent disease or other conditions affecting drug intake which can contribute to the virulence of coccidiosis under field conditions.

Do not use in feeds containing bentonite.

Warning

Keep this and all drugs out of the reach of children.

Manufactured By

Blue Bird Feed Mill
Robin, Indiana 00000

Net Weight __ lbs (__ kg)
Bag or Bulk