

**Blue Bird Bambermycins
(bambermycins Type B medicated feed)**

Supplemental Pasture Cattle Feed

Must Be Mixed Before Feeding

For increased rate of weight gain in pasture cattle (slaughter, stocker and feeder cattle, and dairy and beef replacement heifers).

ACTIVE DRUG INGREDIENT

Bambermycins 100.0 to 800.0 g/ton*

GUARANTEED ANALYSIS

Crude Protein, not less than	_____	%
Non-Protein Nitrogen (NPN) ¹ , not more than	_____	%
Crude Fat, not less than	_____	%
Crude Fiber, not more than	_____	%
Calcium, not less than	_____	%
Calcium, not more than	_____	%
Phosphorus, not less than	_____	%
Salt ² , not less than	_____	%
Salt ² , not more than	_____	%
Sodium ³ , not less than	_____	%
Sodium ³ , not more than	_____	%
Potassium, not less than	_____	%
Vitamin A ^{2,4} , not less than	_____	I.U./lb

¹ When added.

² If added.

³ Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

⁴ Other than precursors of Vitamin A.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

MIXING DIRECTIONS

To manufacture Type C medicated feeds containing 2.0 to 80.0 g/ton bambermycins from Type B medicated feeds mix:

40 to 1600 lbs. of Type B feed containing 100 g/ton bambermycins with 1960 to 400 lbs. of non-medicated feed, or

5 to 200 lbs. of Type B feed containing 800 g/ton bambermycins with 1995 to 1800 lbs. of non-medicated feed.

Thoroughly mix Type B medicated feed containing bambermycins and non-medicated feed to ensure complete and uniform distribution.

Lot No.

MANUFACTURED BY

BLUE BIRD FEED MILL
Any town, USA 12345

Net Weight on Bulk Invoice

* Final printed label on formulated Type B medicated feed must bear a single concentration of each drug.