

Date of Approval: **OCT 26 2007**

**FREEDOM OF INFORMATION SUMMARY**  
**SUPPLEMENTAL NEW ANIMAL DRUG APPLICATION**

NADA 141-246

**AQUAFLO**

**Florfenicol**  
**Type A medicated article**  
**Freshwater-reared salmonids**

**“for the control of mortality in freshwater-reared salmonids due to furunculosis  
associated with *Aeromonas salmonicida*”**

**Sponsored by:**

**Schering-Plough Animal Health Corp.**

**2007-141-246**

**FOIS3**

**TABLE OF CONTENTS**

I. GENERAL INFORMATION:..... 1

II. EFFECTIVENESS:.....2

    A. Dosage Characterization: .....2

    B. Substantial Evidence:.....2

III. TARGET ANIMAL SAFETY:.....7

IV. HUMAN FOOD SAFETY: .....7

    A. Toxicology: .....7

    B. Residue Chemistry:.....7

    C. Microbial Food Safety: .....7

    D. Analytical Method for Residues: .....7

V. USER SAFETY:.....7

VI. AGENCY CONCLUSIONS:..... 8

    A. Marketing Status: .....8

    B. Exclusivity: .....8

    C. Supplemental Applications: .....9

    D. Patent Information: .....9

VII. ATTACHMENTS:.....9

**I. GENERAL INFORMATION:**

- A. File Number:** NADA 141-246
- B. Sponsor:** Schering-Plough Animal Health Corp.  
556 Morris Ave.  
Summit, NJ 07901  
  
Drug Labeler Code: 000061
- C. Proprietary Name(s):** AQUAFLOR
- D. Established Name(s):** Florfenicol
- E. Pharmacological Category:** Antimicrobial
- F. Dosage Form(s):** Type A medicated article
- G. Amount of Active Ingredient(s):** 500 g of florfenicol per kg (227.27 g per lb)
- H. How Supplied:** 2-kg foil laminate foil pouches (12 x 16 inches)  
16-kg fiber board drum (8 x 2-kg pouches)
- I. How Dispensed:** VFD
- J. Dosage(s):** 10 mg florfenicol/kg of fish/day for  
10 consecutive days
- K. Route(s) of Administration:** Oral
- L. Species/Class(es):** Freshwater-reared salmonids
- M. Indication(s):** For the control of mortality due to furunculosis associated with *Aeromonas salmonicida*.
- N. Effect(s) of Supplement:** This supplement provides for the addition of an indication for the control of mortality in freshwater-reared salmonids due to furunculosis associated with *Aeromonas salmonicida*.

## II. EFFECTIVENESS:

### A. Dosage Characterization:

*Aeromonas salmonicida* is the bacterial pathogen associated with furunculosis in freshwater-reared salmonids. This disease is frequently present as a systemic disease that may result in high levels of mortality in affected fish.

The florfenicol dose selected for the effectiveness trials was based on the *A. salmonicida* minimum inhibitory concentration of florfenicol, the pharmacokinetic profile of florfenicol in salmonids, and the results of field studies for saltwater-reared salmonids available in the published literature.

Based on that information, florfenicol administered at a dose of 10 mg/kg of fish daily had the potential to control furunculosis outbreaks associated with *A. salmonicida* in freshwater-reared salmonids.

### B. Substantial Evidence:

The results of the three following studies, when considered together, demonstrate that florfenicol is effective when administered in feed at a dose of 10 mg/kg of fish/day for 10 consecutive days for the control of mortality in freshwater-reared salmonids due to furunculosis associated with *Aeromonas salmonicida*.

#### 1. Field Study

- a. "The Efficacy of Florfenicol-Medicated Feed to Control Mortality of Fingerling Coho Salmon, *Oncorhynchus kisutch*, Caused by Furunculosis, Causative Agent *Aeromonas salmonicida*" (Study Number FLOR-01-EFF-01)
- b. Investigator: Al Jensen  
Makah National Fish Hatchery  
U.S. Fish and Wildlife Service  
Neah Bay, WA
- c. Study Design:
  - 1) Objective: To evaluate the effectiveness of florfenicol administered in feed at a dose of 10 mg florfenicol/kg of fish/day for 10 consecutive days to control mortality in fingerling coho salmon due to furunculosis associated with *A. salmonicida*.
  - 2) Study Animals: Approximately 2,400 fingerling coho salmon
  - 3) Treatment Groups: The study included two treatment groups with six replicates of each treatment. Each replicate was a tank of fish. Treatments were assigned to tanks using a completely randomized study design. One

florfenicol-treated tank was removed from the study because of an interruption in water flow.

- 4) **Drug Administration:** Florfenicol was administered in a commercial salmonid feed at a dose of either 0 or 10 mg florfenicol/kg of fish daily. Study feeds were fed for 10 consecutive days.
- 5) **Measurement and Observations:** *A. salmonicida* was identified on cultures of spleen and kidney tissue collected 5 days before the start of the treatment period (60 fish), and on the first day of the treatment period (54 fish). The clinical signs observed were consistent with a furunculosis infection. Approximately 220 fish were randomly transferred to the study tanks. The study included a one-day acclimation period, a 10-day treatment period, and a 7-day post-treatment period. Dead fish (mortalities) were counted and recorded twice daily. On Day 18, six fish were collected from each tank for examination and collection of samples for culture. Fish examinations on Day 18 revealed an infection with the external parasite, *Trichodina* sp. The study was ended and treatment for the parasite infestation was initiated.
- 6) **Statistical Analysis:** The design of this study precluded statistical analysis.

d. **Results:** Mortality results are included in the following table.

**Table 1.** Mortality results for a field effectiveness study in coho salmon with a 10-day treatment period and 7-day post-treatment period

<b>Florfenicol Dose (mg/kg of fish)</b>	<b>Percent Cumulative Mortality</b>
0	30.3 (331/1,092)
10	11.1 (145/1,306)

- e. **Adverse Reaction:** No adverse reactions were reported in this study.
- f. **Conclusion:** Results of this study support the effectiveness of florfenicol administered in feed at a dose of 10 mg/kg of fish/day for 10 consecutive days to control mortality in coho salmon, *Oncorhynchus kisutch*, due to furunculosis associated with *Aeromonas salmonicida*.

## 2. Field Study

- a. "The Efficacy of Florfenicol-Medicated Feed to Control Mortality of Fall Chinook Salmon, *Oncorhynchus tshawytscha*, Caused by Furunculosis, Causative Agent *Aeromonas salmonicida*" (Study No. FLOR-01-EFF.3-22)

- b. Investigator: Randy Rickert  
Makah National Fish Hatchery  
U.S. Fish and Wildlife Service  
Neah Bay, WA
- c. Study Design:
- 1) Objective: To evaluate the effectiveness of florfenicol administered in feed at a dose of 10 mg florfenicol/kg of fish/day for 10 consecutive days to control mortality in fall chinook salmon fingerlings due to furunculosis associated with *A. salmonicida*.
  - 2) Study Animals: Approximately 1,812 fall chinook salmon fingerlings
  - 3) Treatment Groups: The study included two treatment groups with six replicates of each treatment. Each replicate was a tank of fish. Treatments were assigned to tanks using a randomized block study design.
  - 4) Drug Administration: Florfenicol was administered in a commercial salmonid feed at a dose of either 0 or 10 mg florfenicol/kg of fish daily. Study feeds were fed for 10 consecutive days. The trial feed was assayed to confirm the florfenicol concentration.
  - 5) Measurement and Observations: Fish were held in one raceway prior to allocation to study tanks. Dead, moribund, and healthy fish were collected 5 days prior to the start of the acclimation period for examination. *A. salmonicida* was identified on cultures and imprints of kidney tissue. Equivalent numbers of fish were transferred to each study tank in five rounds, for a total of approximately 151 fish per tank. The study included a one-day acclimation period, a 10-day treatment period, and a 14-day post-treatment period. Dead fish (mortalities) were counted once or twice daily. Dead fish were collected during the treatment period (31 fish) and the post-treatment period (10 fish). Kidney tissue was collected from each fish for culture. *A. salmonicida* was identified via cultures of kidney tissue collected from the fish. Fish behavior and appetite were observed daily.
  - 6) Statistical Analysis: Arcsine-transformed Kaplan Meier estimates of survival were calculated for each tank. Means of treated and control tanks were then compared using a t-test. Significant differences were seen between treated and control groups immediately after treatment, after treatment plus a 10-day holding period, and after treatment plus a 14-day holding period. Mixed model repeated measures analyses using a binomial distribution with canonical logistic link showed that most of the differences in cumulative survival were due to reduced daily mortality rates among treated individuals from 4 to 18 days after the start of florfenicol administration.

- d. Results: Mortality results are included in the following table.

**Table 2.** Mortality results for a field effectiveness study in chinook salmon with a 10-day treatment period and 14-day post-treatment period

<b>Florfenicol Dose (mg/kg of fish)</b>	<b>Percent Cumulative Mortality (Cumulative Mortality)</b>
0	94.4 (808)
10	16.2 (141)

The treated and untreated control groups differ significantly in the cumulative percent mortality ( $p < 0.001$ ).

- e. Adverse Reaction: No adverse reactions were reported in this study.
- f. Conclusion: Results of this study demonstrate the effectiveness of florfenicol administered in feed at a dose of 10 mg/kg of fish/day for 10 consecutive days to control mortality in fingerling fall chinook salmon, *Oncorhynchus tshawytscha*, due to furunculosis associated with *A. salmonicida*.

### 3. Field Study

- a. "The Efficacy of Florfenicol-Medicated Feed to Control Mortality of Coho Salmon, *Oncorhynchus kisutch*, Caused by Furunculosis, Causative Agent *Aeromonas salmonicida*" (Study No. FLOR-01-EFF.3-27)
- b. Investigator: Randy Rickert  
Makah National Fish Hatchery  
U.S. Fish and Wildlife Service  
Neah Bay, WA
- c. Study Design:
- 1) **Objective**: To evaluate the effectiveness of florfenicol administered in feed at a dose of 10 mg florfenicol/kg of fish/day for 10 consecutive days to control mortality in coho salmon fingerlings due to furunculosis associated with *A. salmonicida*.
  - 2) **Study Animals**: Approximately 1,920 coho salmon fingerlings
  - 3) **Treatment Groups**: The study included two treatment groups with six replicates of each treatment. Each replicate was a tank of fish. Treatments were assigned to tanks using a randomized block study design.
  - 4) **Drug Administration**: Florfenicol was administered in a commercial salmonid feed at a dose of either 0 or 10 mg florfenicol/kg of fish daily. Study feeds were fed for 10 consecutive days. The trial feed was assayed to confirm the florfenicol concentration.

- 5) **Measurement and Observation:** Fish were held in one raceway prior to allocation to experimental tanks. Dead and moribund fish were collected 2 days prior to the start of the acclimation period for examination and tissue collection for culture. *A. salmonicida* was identified on cultures of kidney tissue. Equivalent numbers of fish were randomly transferred to each study tank in four rounds, for a total of approximately 160 fish per tank. The study included a 2-day acclimation period, a 10-day treatment period, and a 14-day post-treatment period. Dead fish (mortalities) were counted once or twice daily. Moribund fish were collected during the treatment period (3 fish) and the post-treatment period (2 fish). Kidney tissue was collected from each fish for culture. *A. salmonicida* was identified via cultures of kidney tissue collected from control group fish. Fish behavior and appetite were observed daily.
- 6) **Statistical Analysis:** Arcsine-transformed proportion survival was calculated for each tank. Means of treated and control tanks were then compared using a t-test. Significant differences were seen between treated and control groups immediately after treatment, after treatment plus a 10-day holding period, and after treatment plus a 14-day holding period. Mixed model repeated measures analyses on arcsine transformed daily mortality rates showed that most of the differences in cumulative survival were due to reduced daily mortality rates among treated individuals from 8 to 19 days after the start of florfenicol administration.

d. **Results:** Mortality results are included in the following table.

**Table 3.** Mortality results for a field effectiveness study in coho salmon with a 10-day treatment period and 7-day post-treatment period.

<b>Florfenicol Dose (mg/kg of fish)</b>	<b>Percent Cumulative Mortality (Cumulative Mortality)</b>
0	29.5 (282/956)
10	16.0 (151/946)

The treated and untreated control groups differ significantly in the cumulative percent mortality ( $p < 0.005$ ).

- e. **Adverse Reactions:** No adverse reactions were reported in this study.
- f. **Conclusion:** Results of this study demonstrate the effectiveness of florfenicol administered in feed at a dose of 10 mg/kg of fish/day for 10 consecutive days to control mortality in fingerling coho salmon, *Oncorhynchus kisutch*, due to furunculosis associated with *A. salmonicida*.

### **III. TARGET ANIMAL SAFETY:**

CVM did not require target animal safety studies for this supplemental approval. The FOI Summary for the supplemental approval of NADA 141-246 dated March 19, 2007, contains a summary of target animal safety studies for the use of florfenicol at a dose of 10 mg/kg of fish/day for 10 consecutive days in freshwater-reared salmonids.

### **IV. HUMAN FOOD SAFETY:**

#### **A. Toxicology:**

CVM did not require toxicology studies for this supplemental approval. The FOI Summary for the original approval of NADA 141-063 (florfenicol in cattle) dated May 31, 1996, and for the supplemental approval of NADA 141-246 dated March 19, 2007, contain summaries of all toxicology studies. The ADI for florfenicol is 10 micrograms per kilogram body weight per day. The safe concentration of total drug-related residues is 2 ppm in salmonid muscle/skin.

#### **B. Residue Chemistry:**

CVM did not require residue chemistry studies for this supplemental approval. The FOI Summary for the supplemental approval of NADA 141-246 dated March 19, 2007, contains a summary of residue chemistry studies for freshwater-reared salmonids.

#### **C. Microbial Food Safety:**

CVM considered the impact of adding a claim for the control of mortality due to furunculosis to the approved label for florfenicol for freshwater-reared salmonids on microbial food safety, and determined that additional microbial food safety information for this supplemental approval was not required. The FOI Summary for the supplemental approval of NADA 141-246, dated March 19, 2007, contains microbial food safety information regarding the use of florfenicol in freshwater-reared salmonids.

#### **D. Analytical Method for Residues:**

The FOI Summary for the supplemental approval of NADA 141-246 dated March 19, 2007, contains the analytical method summaries for florfenicol in salmonids.

### **V. USER SAFETY:**

The product labeling contains the following information regarding safety to humans handling, administering, or exposed to AQUAFLO<sup>R</sup>:

“Avoid inhalation, oral exposure, and direct contact with skin or eyes. Operators mixing and handling AQUAFLOR (florfenicol) should use protective clothing, gloves, goggles and NIOSH-approved dust mask. Wash thoroughly with soap and water after handling. If accidental eye contact occurs, immediately rinse thoroughly with water. If irritation persists, seek medical attention. Not for human consumption. Keep out of reach of children. The Material Safety Data Sheet (MSDS) contains more detailed occupational safety information. For a copy of the MSDS sheet, call 1-800-770-8878. For more information or to report adverse effects, call 1-800-211-3573.”

CVM examined the material safety data sheet to conclude that user safety concerns have been appropriately addressed in the labeling.

## **VI. AGENCY CONCLUSIONS:**

The data submitted in support of this NADA satisfy the requirements of section 512 of the Federal Food, Drug, and Cosmetic Act and 21 CFR Part 514. The data demonstrate that AQUAFLOR, when used according to the label, is safe and effective for the control of mortality due to furunculosis associated with *Aeromonas salmonicida*. Additionally, data demonstrate that residues in food products derived from freshwater-reared salmonids treated with AQUAFLOR will not represent a public health concern when the product is used according to the label.

### **A. Marketing Status:**

This drug may be dispensed only under a valid Veterinary Feed Directive (VFD). Any animal feed bearing or containing this VFD drug will be fed to animals only by or on a lawful veterinary feed directive issued by a licensed veterinarian in the course of the veterinarian’s professional practice. In addition, veterinary feed directives issued for this drug are not refillable.

Labeling restricts this drug to use by or on the order of a licensed veterinarian. The decision to restrict this drug to use by or on the order of a licensed veterinarian was based on the following factors: (a) adequate directions cannot be written to enable lay persons to appropriately diagnose and subsequently use this product, (b) restricting this drug to use by or on the order of a licensed veterinarian should help prevent indiscriminate use which could result in violative tissue residues, and (c) the rate of emergence of florfenicol-resistant organisms may be reduced by the involvement of veterinarians in product use. Because the drug will be administered in feed, the drug will be marketed as a VFD drug.

### **B. Exclusivity:**

AQUAFLOR in the dosage form and for the intended uses approved by FDA, for the control of mortality in freshwater-reared salmonids due to furunculosis associated with *Aeromonas salmonicida*, qualifies for seven years of exclusive marketing rights beginning as of the date of approval. This new animal drug qualifies for exclusive

marketing rights under section 573(c) of the Federal Food, Drug, and Cosmetic Act (the act) because it has been declared a designated new animal drug by FDA under section 573(a) of the act.

**C. Supplemental Applications:**

This supplemental NADA did not require a reevaluation of the safety or effectiveness data in the original NADA [21 CFR §514.106(b)(2)].

**D. Patent Information:**

The sponsor did not submit any patent information with this application.

**VII. ATTACHMENTS:**

Facsimile Labeling:

AQUAFLO (florfenicol) Type A medicated article label 2 kg

AQUAFLO (florfenicol) Type A medicated article label 8 x 2 kg

AQUAFLO (florfenicol) Type C medicated feed label for fresh-water reared salmonids

AQUAFLO (florfenicol) VFD Form

NDC XXXX-XXXX-XX

2.0 kg (4.4 lb)

# Aquaflor<sup>®</sup>

(Florfenicol)

Type A Medicated Article  
For Use in Catfish and Salmonid Feeds Only

**Do Not Feed Undiluted**

Active Drug Ingredient: Florfenicol 500 g per kg (227.27 g per lb)

Description: Each kg of Aquaflor (Florfenicol) contains 500 g (1.1 lb) of florfenicol in a palatable base.

NADA #141-246, Approved by FDA



Schering-Plough

Copyright © 2005-2007, Schering-Plough Animal Health Corporation, Summit, NJ 07901, USA. Made in Ireland. All rights reserved.

XXXXXXXX Rev. 10/07

NDC XXXX-XXXX-XX

2.0 kg (4.4 lb)

# Aquaflor® (Florfenicol)

Type A Medicated Article  
For Use in Catfish and Salmonid Feeds Only

Do Not Feed Undiluted

**CAUTION:** Federal law limits this drug to use under the professional supervision of a licensed veterinarian. Animal feed bearing or containing this veterinary feed directive drug shall be fed to animals only by or upon a lawful veterinary feed directive (VFD) issued by a licensed veterinarian in the course of the veterinarian's professional practice.

**Active Drug Ingredient:** Florfenicol 500 g per kg (227.27 g per lb)  
**Inert ingredients:** Lactose and Povidone.

**Description:** Each kg of Aquaflor® (florfenicol) contains 500 g (1.1 lb) of florfenicol in a palatable base.

**Activity:**

**Catfish:** *In vitro* and *in vivo* investigations in catfish have established florfenicol's activity against *Edwardsiella ictaluri* (Table 1).

Table 1: Minimum inhibitory concentration (MIC) of florfenicol against *Edwardsiella ictaluri* isolated from channel catfish, between 1998-2001.

Organism	No. of Isolates	MIC <sub>50</sub> (µg/mL)	MIC <sub>90</sub> (µg/mL)
<i>Edwardsiella ictaluri</i>	95	0.25	0.25

**Indications:**

**Catfish:** For the control of mortality in catfish due to enteric septicemia of catfish associated with *Edwardsiella ictaluri*.

**Freshwater-reared Salmonids:** For the control of mortality in freshwater-reared salmonids due to coldwater disease associated with *Flavobacterium psychrophilum* and furunculosis associated with *Aeromonas salmonicida*.

**Caution:** The effects of Aquaflor® (florfenicol) on reproductive performance have not been determined. For catfish, a dose-related decrease in hematopoietic/lymphopoietic tissue may occur. The time required for the hematopoietic/lymphopoietic tissues to regenerate was not evaluated.

▶ **RESIDUE WARNING:** Catfish: Feeds containing Aquaflor® (florfenicol) must be withdrawn 12 days prior to slaughter.

▶ **RESIDUE WARNING:** Freshwater-reared Salmonids: Feeds containing Aquaflor® (florfenicol) must be withdrawn 15 days prior to slaughter.

**IMPORTANT:** Must be thoroughly mixed in feeds or surface-coated (top-coated) onto the feeds before use.

**Mixing Instructions:**

**For incorporation inside pellets:** For making Aquaflor® (florfenicol) Type C Medicated Feed for catfish and freshwater-reared salmonids: a) Aquaflor® (florfenicol) is added to other feed ingredients in the mixer, b) the medicated feed is mixed thoroughly to insure homogeneity, c) the mixture is extruded and pellets are dried, d) the pellets are dry-mixed or coated with a predetermined amount of fish or vegetable oil, and e) at the completion of mixing, the product is transferred to a storage tank for packaging or transport.

**For surface-coating (top-coating):** For making Aquaflor® (florfenicol) Type C Medicated Feed for freshwater-reared salmonids only: a) add a known quantity of fish feed into a mixer, b) weigh out Aquaflor® (florfenicol), c) weigh out fish oil or vegetable oil into a bucket, d) mix Aquaflor® (florfenicol) and oil thoroughly in the bucket, e) add the Aquaflor® (florfenicol) and oil mixture

to the feed in the mixer, slowly, while the mixer is running at low speed, f) at the completion of mixing, the product is transferred to a storage tank for packaging or transport.

Recommended Aquaflor® (florfenicol) Inclusion Rates for Preparation of Type C Medicated Feed for Catfish and Freshwater-reared Salmonids

Feeding Rate	Florfenicol Concentration in Feed	Amount of Aquaflor® (florfenicol) per Ton	Biomass of Fish Medicated per Ton of Feed per 10-day Treatment Period
	Grams/lb	lbs	lbs
0.5	1,816	8.00	40,000
1.0	908	4.00	20,000
2.0	454	2.00	10,000
3.0	300	1.32	6,666
5.0	182	0.80	4,000

**Feeding Directions:**

**Catfish:** Feed as the sole ration for 10 consecutive days. Aquaflor® (florfenicol) medicated feed should only be administered once disease associated with *Edwardsiella ictaluri* in catfish has been appropriately diagnosed. Feeding fish at a percent of biomass and corresponding florfenicol concentration included in the table above will deliver 10 mg florfenicol per kg of fish.

**Freshwater-reared salmonids:** Feed as the sole ration for 10 consecutive days. Aquaflor® (florfenicol) medicated feed should only be administered once disease associated with *Flavobacterium psychrophilum* or *Aeromonas salmonicida* in freshwater-reared salmonids has been appropriately diagnosed. Feeding fish at a percent of biomass and corresponding florfenicol concentration included in the table above will deliver 10 mg florfenicol per kg of fish.

**Caution:** Feed containing Aquaflor® (florfenicol) shall not be fed to catfish or freshwater-reared salmonids for more than 10 days. Following administration, fish should be reevaluated by a licensed veterinarian before initiating a further course of therapy. The expiration date for VFD for Aquaflor® (florfenicol) must not exceed 15 days from the date of issuance. VFD for Aquaflor® (florfenicol) shall not be refilled.

**WARNING:** Avoid inhalation, oral exposure, and direct contact with skin or eyes. Operators mixing and handling Aquaflor® (florfenicol) should use protective clothing, gloves, goggles and NIOSH-approved dust mask. Wash thoroughly with soap and water after handling. If accidental eye contact occurs, immediately rinse thoroughly with water. If irritation persists, seek medical attention. Not for human consumption. Keep out of reach of children. The Material Safety Data Sheet (MSDS) contains more detailed occupational safety information. For a copy of MSDS sheet, call 1-800-770-8878. For more information or to report adverse effects, call 1-800-211-3573.

For customer service, call 1-800-521-5757.

**STORAGE CONDITIONS:** Store at 2°-30°C (36°-86°F).

NADA #141-246, Approved by FDA.

Aquaflor® is a registered trademark of Schering-Plough Animal Health Corporation.

 Schering-Plough

Copyright © 2005, 2007, Schering-Plough Animal Health Corporation, Summit, NJ 07901, USA.  
Made in Ireland. All rights reserved. XXXXXXXX Rev. 10/07



NDC XXXX-XXXX-XX

8 x 2.0 kg  
16 kg (35.2 lb)

# Aquaflor<sup>®</sup>

(Florfenicol)

Type A Medicated Article  
For Use in Catfish and Salmonid Feeds Only

**Do Not Feed Undiluted**

**Active Drug Ingredient:** Florfenicol 500 g per kg (227.27 g per lb)

**Description:** Each kg of Aquaflor<sup>®</sup> (Florfenicol) contains 500 g (1.1 lb) of florfenicol in a palatable base.

NADA #141-246, Approved by FDA.



Schering-Plough

Copyright © 2005, 2007, Schering-Plough Animal Health Corporation, Summit, NJ 07901, USA. Made in Ireland. All rights reserved.

XXXXXXXX Rev. 10/07

NDC XXXX-XXXX-XX

8 x 2.0 kg  
16 kg (35.2 lb)

# Aquaflor® (Florfenicol)

Type A Medicated Article  
For Use in Catfish and Salmonid Feeds Only

**Do Not Feed Undiluted**

**CAUTION:** Federal law limits this drug to use under the professional supervision of a licensed veterinarian. Animal feed bearing or containing this veterinary feed directive drug shall be fed to animals only by or upon a lawful veterinary feed directive (VFD) issued by a licensed veterinarian in the course of the veterinarian's professional practice.

**Active Drug Ingredient:** Florfenicol 500 g per kg (227.27 g per lb)  
**Inert Ingredients:** Lactose and Povidone.

**Description:** Each kg of Aquaflor® (florfenicol) contains 500 g (1.1 lb) of florfenicol in a palatable base.

**Activity:**

**Catfish:** *In vitro* and *in vivo* investigations in catfish have established florfenicol's activity against *Edwardsiella ictaluri* (Table 1).

**Table 1:** Minimum inhibitory concentration (MIC) of florfenicol against *Edwardsiella ictaluri* isolated from channel catfish, between 1998-2001.

Organism	No. of Isolates	MIC <sub>50</sub> (ppm)	MIC Range (ppm)
<i>Edwardsiella ictaluri</i>	95	0.25	0.25

**Indications:**

**Catfish:** For the control of mortality in catfish due to enteric septicemia of catfish associated with *Edwardsiella ictaluri*.

**Freshwater-reared Salmonids:** For the control of mortality in freshwater-reared salmonids due to coldwater disease associated with *Flavobacterium psychrophilum* and furunculosis associated with *Aeromonas salmonicida*.

**Caution:** The effects of Aquaflor® (florfenicol) on reproductive performance have not been determined. For catfish, a dose-related decrease in hematopoietic/lymphopoietic tissue may occur. The time required for the hematopoietic/lymphopoietic tissues to regenerate was not evaluated.

▶ **RESIDUE WARNING:** Catfish: Feeds containing Aquaflor® (florfenicol) must be withdrawn 12 days prior to slaughter. ◀

▶ **RESIDUE WARNING:** Freshwater-reared Salmonids: Feeds containing Aquaflor® (florfenicol) must be withdrawn 15 days prior to slaughter. ◀

**IMPORTANT:** Must be thoroughly mixed in feeds or surface-coated (top-coated) onto the feeds before use.

**Mixing Instructions:**

**For incorporation inside pellets:** For making Aquaflor® (florfenicol) Type C Medicated Feed for catfish and freshwater-reared salmonids: a) Aquaflor® (florfenicol) is added to other feed ingredients in the mixer, b) the medicated feed is mixed thoroughly to insure homogeneity, c) the mixture is extruded and pellets are dried, d) the pellets are dry-mixed or coated with a predetermined amount of fish or vegetable oil, and e) at the completion of mixing, the product is transferred to a storage tank for packaging or transport.

**For surface-coating (top-coating):** For making Aquaflor® (florfenicol) Type C Medicated Feed for freshwater-reared salmonids only: a) add a known quantity of fish feed into a mixer, b) weigh out Aquaflor® (florfenicol), c) weigh out fish oil or vegetable oil into a bucket, d) mix Aquaflor® (florfenicol) and oil thoroughly in the bucket, e) add the Aquaflor® (florfenicol) and oil mixture to

the feed in the mixer, slowly, while the mixer is running at low speed, f) at the completion of mixing, the product is transferred to a storage tank for packaging or transport.

**Recommended Aquaflor® (florfenicol) Inclusion Rates for Preparation of Type C Medicated Feed for Catfish and Freshwater-reared Salmonids**

Feeding Rate	Florfenicol Concentration in Feed	Amount of Aquaflor® (florfenicol) per Ton	Biomass of Fish Medicated per Ton of Feed per 10-day Treatment Period
% Biomass	Grams/ton	lbs	lbs
0.5	1,816	8.00	40,000
1.0	908	4.00	20,000
2.0	454	2.00	10,000
3.0	300	1.32	6,666
5.0	182	0.80	4,000

**Feeding Directions:**

**Catfish:** Feed as the sole ration for 10 consecutive days. Aquaflor® (florfenicol) medicated feed should only be administered once disease associated with *Edwardsiella ictaluri* in catfish has been appropriately diagnosed. Feeding fish at a percent of biomass and corresponding florfenicol concentration included in the table above will deliver 10 mg florfenicol per kg of fish.

**Freshwater-reared salmonids:** Feed as the sole ration for 10 consecutive days. Aquaflor® (florfenicol) medicated feed should only be administered once disease associated with *Flavobacterium psychrophilum* or *Aeromonas salmonicida* in freshwater-reared salmonids has been appropriately diagnosed. Feeding fish at a percent of biomass and corresponding florfenicol concentration included in the table above will deliver 10 mg florfenicol per kg of fish.

**Caution:** Feed containing Aquaflor® (florfenicol) shall not be fed to catfish or freshwater-reared salmonids for more than 10 days. Following administration, fish should be reevaluated by a licensed veterinarian before initiating a further course of therapy. The expiration date for VFD for Aquaflor® (florfenicol) must not exceed 15 days from the date of issuance. VFD for Aquaflor® (florfenicol) shall not be refilled.

**WARNING:** Avoid inhalation, oral exposure, and direct contact with skin or eyes. Operators mixing and handling Aquaflor® (florfenicol) should use protective clothing, gloves, goggles and NIOSH-approved dust mask. Wash thoroughly with soap and water after handling. If accidental eye contact occurs, immediately rinse thoroughly with water. If irritation persists, seek medical attention. Not for human consumption. Keep out of reach of children. The Material Safety Data Sheet (MSDS) contains more detailed occupational safety information. For a copy of MSDS sheet, call 1-800-770-8878. For more information or to report adverse effects, call 1-800-211-3573.

For customer service, call 1-800-521-5767.

**STORAGE CONDITIONS:** Store at 2°-30°C (36°-86°F).

NADA #141-246, Approved by FDA.

Aquaflor® is a registered trademark of Schering-Plough Animal Health Corporation.

 Schering-Plough

Copyright © 2005, 2007, Schering-Plough Animal Health Corporation, Summit, NJ 07901, USA.  
Made in Ireland. All rights reserved. XXXXXXXX Rev. 10/07



**BLUEBIRD FEED COMPANY  
BLUEBIRD SALMONID FEED  
Medicated Type C Feed**

**FOR USE IN FRESHWATER- REARED SALMONIDS ONLY**

**Indication**

For the control of mortality in freshwater-reared salmonids due to coldwater disease associated with *Flavobacterium psychrophilum* and furunculosis associated with *Aeromonas salmonicida*.

**Active Drug Ingredient**

Florfenicol..... 182 to 1816 grams per ton\*

**Guaranteed Analysis**

Crude Protein (min).....%  
Crude Fat (min).....%  
Crude Fiber (max).....%  
Phosphorus (min).....%

**Ingredients**

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials (AAFCO).

**Feeding Directions**

Feed as a sole ration at a rate of \_\_\_\_\_% biomass daily for 10 consecutive days. Feeding at this rate will deliver 10 mg florfenicol per kg of fish\*\*.

**Caution**

Federal law limits this drug to use under the professional supervision of a licensed veterinarian. Animal feed bearing or containing this veterinary feed directive (VFD) drug shall be fed to animals only by or upon a lawful veterinary feed directive issued by a licensed veterinarian in the course of the veterinarian's professional practice.

Feed containing Aquaflor® (florfenicol) shall not be fed to freshwater-reared salmonids for more than 10 days. Following 10 days administration, fish should be reevaluated by a licensed veterinarian before initiating a further course of therapy. The expiration date for VFD for Aquaflor® (florfenicol) must not exceed 15 days from the date of issuance. VFD for Aquaflor® (florfenicol) shall not be refilled.

The effects of florfenicol on reproductive performance have not been determined.

**WARNING: Feeds containing Aquaflor®  
(florfenicol) must be withdrawn 15 days prior to  
slaughter.**

**Storage Conditions:** Store in a cool, dry place. Avoid excessive moisture and heat.

**Manufactured By:** Bluebird Feed Mill, Robin, IN 46813

NET WEIGHT 50 lbs (22.7 kg)

2/07

Feed Lot No. \_\_\_\_\_

Feed Manufacturing Date \_\_\_\_\_

Aquaflor® is a registered trademark of Schering-Plough Animal Health Corporation.

Schering-Plough Animal Health Corporation, Summit, NJ 07901, USA.

Made in Ireland. All rights reserved. 10/07

\*Final printed label must bear a single drug concentration.

\*\*Feed according to the veterinarian instructions on the VFD.



# Aquaflor<sup>®</sup> Veterinary Feed Directive (Florfenicol)



Client: \_\_\_\_\_  
 Address: \_\_\_\_\_  
 \_\_\_\_\_  
 Phone: \_\_\_\_\_  
 Fax: \_\_\_\_\_  
 E-mail: \_\_\_\_\_

Veterinarian: \_\_\_\_\_  
 Address: \_\_\_\_\_  
 \_\_\_\_\_  
 Phone: \_\_\_\_\_  
 Fax: \_\_\_\_\_  
 E-mail: \_\_\_\_\_

**Catfish and/or Freshwater-reared Salmonids to be Treated:** Number, Total Weight (Biomass):  
**Farm Location:** Farm Address, Raceway/Pond Identification (Raceway/Pond Number, etc.):

**Indications:** Catfish: For the control of mortality in catfish due to enteric septicemia of catfish associated with *Edwardsiella ictaluri*.  
 Freshwater-reared Salmonids: For the control of mortality in freshwater-reared salmonids due to coldwater disease associated with *Flavobacterium psychrophilum* and furunculosis associated with *Aeromonas salmonicida*.

**Mix into Type C Medicated Feed to Provide:** Check one:  182 g/ton     300 g/ton     454 g/ton     908 g/ton     1816 g/ton

**Amount of Final (Type C) Feed:** \_\_\_\_\_ (Pounds or Tons)    **Feeding Rate:** \_\_\_\_\_ % Biomass

**Feeding Directions:** Feed as the sole ration for 10 consecutive days. Feeding at this rate will deliver 10 mg florfenicol per kg of fish.

Feeding Rate	Florfenicol Concentration in Feed	Amount of Aquaflor <sup>®</sup> (florfenicol) per Ton	Biomass of Fish Medicated per Ton of Feed per 10-day Treatment Period
% Biomass	Grams/ton	lbs	lbs
0.5	1,816	8.00	40,000
1.0	908	4.00	20,000
2.0	454	2.00	10,000
3.0	300	1.32	6,666
5.0	182	0.80	4,000

**Special Instructions**

**Date of Treatment:** \_\_\_\_\_ (Month/Day/Year)

**Expiration Date of VFD:** \_\_\_\_\_ Month/Day/Year (Not to exceed 15 days from date of issuance.)

**Veterinarian's Signature:** \_\_\_\_\_ Date: \_\_\_\_\_

**License Number and State:** \_\_\_\_\_

**Caution:** Federal law limits this drug to use under the professional supervision of a licensed veterinarian. Animal feed bearing or containing this veterinary feed directive drug shall be fed to animals only by or upon a lawful veterinary feed directive (VFD) issued by a licensed veterinarian in the course of the veterinarian's professional practice. Extra-label use (i.e., use of this VFD feed in a manner other than as provided by the VFD drug approval) is strictly prohibited.

Feed containing Aquaflor<sup>®</sup> (florfenicol) shall not be fed to catfish or freshwater-reared salmonids for more than 10 days. Following 10 days administration, fish should be re-evaluated by a licensed veterinarian before initiating a further course of therapy. The expiration date for VFD for Aquaflor (florfenicol) must not exceed 15 days from the date of issuance. VFD for Aquaflor (florfenicol) shall not be refilled. For catfish, a dose related decrease in hematopoietic/lymphopoietic tissue may occur. The time required for the hematopoietic/lymphopoietic tissues to regenerate was not evaluated. The effects of florfenicol on reproductive performance have not been determined.

**RESIDUE WARNING:** Catfish: Feeds containing Aquaflor<sup>®</sup> (florfenicol) must be withdrawn 12 days prior to slaughter.

**RESIDUE WARNING:** Freshwater-reared Salmonids: Feeds containing Aquaflor<sup>®</sup> (florfenicol) must be withdrawn 15 days prior to slaughter.

**Warning:** Avoid inhalation, oral exposure, and direct contact with skin or eyes. Operators mixing and handling Aquaflor<sup>®</sup> should use protective clothing, gloves, goggles and NIOSH-approved dust mask. Wash thoroughly with soap and water after handling. If accidental eye contact occurs, immediately rinse thoroughly with water. If irritation persists, seek medical attention. Not for human consumption. Keep out of reach of children. The Material Safety Data Sheet (MSDS) contains more detailed occupational safety information. For a copy of the MSDS, call 1-800-770-8878. To obtain more information or to report adverse effects, call 1-800-211-3573.

