

# **Exhibit I**

**Label for Paylean™**

# ELANCO

AF0602-25B

**For Use in Finishing  
Swine Feeds Only**

# Paylean<sup>®</sup> 9

**Ractopamine  
Hydrochloride**

**Net Weight 25 lb  
(11.34 kg)**

## Type A Medicated Article

**Do not feed undiluted.**

**Active Drug Ingredient:** ractopamine hydrochloride – 9 g per lb (20 g per kg)

**Important:** Must be thoroughly mixed into feeds before use. Follow label directions.

**Indication:** For increased rate of weight gain, improved feed efficiency, and increased carcass leanness in finishing swine fed a complete ration containing at least 16% crude protein from 150 lb (68 kg) to 240 lb (109 kg) body weight.

Indications	Appropriate Concentration of Ractopamine in Type C Medicated Feed
Increased Rate of Weight Gain, Improved Feed Efficiency, and Increased Carcass Leanness	4.5 grams/ton (5 ppm)
Improved Feed Efficiency and Increased Carcass Leanness	4.5 to 18 grams/ton (5 ppm to 20 ppm)

**Inert Ingredients:** Ground corncobs.

Carcass Measurements	Effect of Ractopamine	
	4.5 grams/ton (5 ppm)	9 - 18 grams/ton (10 - 20 ppm)
Carcass Fat	NC	↓
10th Rib Backfat (3/4 location)	NC	↓
Last Rib Backfat (midline)	NC	NC
Loineye Area (10th rib)	NC	↑
Rate of Lean Gain	NC	↑
Efficiency of Lean Gain	NC	↑
Dressing Percentage	NC	↑

NC= No Change, ↑ = increased, ↓ = decreased

**Mixing Directions:** Thoroughly mix Paylean 9 Type A Medicated Article into one ton of appropriate feed ingredients or diluents according to the table below to obtain the proper concentration in the Type B Medicated Feed (maximum 3600 g/ton). The following table gives examples of how some Type B Medicated Feed concentrations can be prepared:

Pounds of Paylean 9 To Add Per Ton To Make a Type B Medicated Feed	Resulting Ractopamine Concentration in Type B Medicated Feed	
	grams/ton	grams/pound
100	900	0.45
200	1,800	0.90
300	2,700	1.35
400	3,600	1.80

Thoroughly mix Paylean 9 Type A Medicated Article into one ton of complete swine feed according to the table below to obtain the proper concentration in the Type C Medicated Feed. Prepare an intermediate pre-blend of the premix prior to mixing in a complete feed. Thoroughly mix the required amount in a convenient quantity of feed ingredients then add to the remaining feed ingredients to make a ton of complete feed.

Pounds Paylean 9 To Add Per Ton of Type C Medicated Feed	Resulting Ractopamine Concentration in Type C Medicated Feed	
0.5	4.5	grams/ton ( 5 ppm)
1.0	9	grams/ton (10 ppm)
1.5	13.5	grams/ton (15 ppm)
2.0	18	grams/ton (20 ppm)

**Feeding Directions:** Feed continuously to finishing swine as the sole ration from 150 lb (68 kg) to 240 lb (109 kg) body weight.

**CAUTION:** Not for use in breeding swine.

**WARNING:** The active ingredient in Paylean, ractopamine hydrochloride, is a beta-adrenergic agonist. Individuals with cardiovascular disease should exercise special caution to avoid exposure. Not for use in humans. Keep out of the reach of children. The Paylean 9 formulation (Type A Medicated Article) poses a low dust potential under usual conditions of handling and mixing. When mixing and handling Paylean, use protective clothing, impervious gloves, protective eye wear, and a NIOSH-approved dust mask. Operators should wash thoroughly with soap and water after handling. If accidental eye contact occurs, immediately rinse eyes thoroughly with water. If irritation persists, seek medical attention. The material safety data sheet contains more detailed occupational safety information. To report adverse effects, access medical information, or obtain additional product information, call 1-800-428-4441.

Store at room temperature.

Expiration Date and Lot Number are printed on the bag. Not to be used after the expiry date.

## Paylean® 9

Elanco Animal Health  
A Division of Eli Lilly and Company  
Indianapolis, IN 46285, U.S.A.



Questions or Comments: Call 1-800-428-4441

Paylean® is a registered trademark of Eli Lilly and Company