

90000 SERIES
30% P.C.W.


§ 558.76

21 CFR Ch. I (4-1-03 Edition)

(2) Arsanilic acid may be used in accordance with the provisions of this section in the combinations provided as follows:

- (i) Amprolium in accordance with § 558.55.
- (ii) Amprolium and ethopabate in accordance with § 558.58.
- (iii) Bacitracin zinc in accordance with § 558.78.
- (iv) Bacitracin and zoalene in accordance with § 558.680.
- (v) Zoalene in accordance with § 558.680.

[41 FR 10992, Mar. 15, 1976, as amended at 42 FR 18617, Apr. 8, 1977; 51 FR 7395, Mar. 3, 1986; 51 FR 33897, Sept. 24, 1986; 54 FR 18281, Apr. 28, 1989; 56 FR 19268, Apr. 26, 1991; 60 FR 39847, Aug. 4, 1995; 66 FR 14074, Mar. 9, 2001; 66 FR 57873, Nov. 19, 2001]

§ 558.76 Bacitracin methylene disalicylate.

(a) *Approvals.* Type A medicated articles: 10, 25, 30, 40, 50, 60, or 75 grams per pound to 046573 in § 510.600(c) of this chapter.

(b) *Special considerations.* The quantities of antibiotics are expressed in terms of the equivalent amount of antibiotic standard.

(c) *Related tolerances.* See § 556.70 of this chapter.

(d) *Conditions of use.* (1) It is used as follows:

Bacitracin methylene disalicylate in grams per ton	Combination in grams per ton	Indications for use	Limitations	Sponsor
(i) 4 to 50		Chickens, turkeys, and pheasants; increased rate of weight gain and improved feed efficiency ¹ .		046573
(ii) 5 to 20		Quail not over 5 weeks of age, increased rate of weight gain and improved feed efficiency ¹ .		046573
(iii) 10 to 25		Chickens; for increased egg production and improved feed efficiency for egg production	For first 7 months of production	046573
(iv) 10 to 30		Swine: for increased rate of weight gain and improved feed efficiency.	For growing and finishing swine	046573
	Chlortetracycline approximately 400, varying with body weight and food consumption to provide 10 milligrams per pound of body weight per day.	Swine; for increased rate of weight gain and improved feed efficiency; for treatment of bacterial enteritis caused by <i>Escherichia coli</i> and <i>Salmonella choleraesuis</i> and bacterial pneumonia caused by <i>Pasteurella multocida</i> susceptible to chlortetracycline.	Feed for not more than 14 days to provide 10 milligrams of chlortetracycline per pound of body weight per day; as chlortetracycline provided by No. 046573 in § 510.600(c) of this chapter. Type C feed may be prepared from Type B feed containing 1 to 3 grams per pound BMD with 400 grams per pound CTC, to 046573 in § 510.600(c).	046573
		Swine, for control of porcine proliferative enteropathies (ileitis) caused by <i>Lawsonia intracellularis</i> susceptible to chlortetracycline.	Feed for not more than 14 days, chlortetracycline and BMD® as provided by 046573 in § 510.600(c) of this chapter.	046573
(v) [Reserved]				
(vi) 50		Broiler chickens; as an aid in the prevention of necrotic enteritis caused or complicated by <i>Clostridium</i> spp. or other organisms susceptible to bacitracin.		046573
		Replacement chickens; as an aid in the prevention of necrotic enteritis caused or complicated by <i>Clostridium</i> spp. or other organisms susceptible to bacitracin	Feed continuously as sole ration	046573

Bacitracin methylene disalicylate in grams per ton	Combination in grams per ton	Indications for use	Limitations	Sponsor
(vii)-(viii) [Reserved]				
(ix) 100 to 200		Broiler chickens; as an aid in the control of necrotic enteritis caused or complicated by <i>Clostridium</i> spp. or other organisms susceptible to bacitracin.		046573
		Replacement chickens; as an aid in the control of necrotic enteritis caused or complicated by <i>Clostridium</i> spp. or other organisms susceptible to bacitracin	Feed continuously as sole ration. Start at first clinical signs of disease, vary dosage based on severity of infection, administer continuously for 5 to 7 days or as long as clinical signs persist, then reduce medication to prevention level (50 g/t).	046573
(x) 200		Turkeys; as an aid in the control of transmissible enteritis in growing turkeys complicated by organisms susceptible to bacitracin methylene disalicylate.		046573
		Quail; for the prevention of ulcerative enteritis in growing quail due to <i>Clostridium colinum</i> susceptible to bacitracin methylene disalicylate	From Type A medicated articles containing 25, 40, or 50 grams of bacitracin methylene disalicylate. Feed continuously as the sole ration.	046573
(xi) 250		1. Growing/Finishing Swine: For control of swine dysentery associated with <i>Treponema hyodysenteriae</i> on premises with a history of swine dysentery but where signs of the disease have not yet occurred; or following an approved treatment of the disease condition	As the sole ration. Not for use in swine weighing more than 250 pounds. Diagnosis should be confirmed by a veterinarian when results are not satisfactory.	046573
		2. Pregnant sows: For control of clostridial enteritis caused by <i>C. perfringens</i> in suckling piglets.	As the sole ration. Feed to sows from 14 days before through 21 days after farrowing on premises with a history of clostridial scours. Diagnosis should be confirmed by a veterinarian when results are not satisfactory	

¹ These conditions are NAS/NRC reviewed and found effective. Applications for these uses may not require effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

(2) It is used as bacitracin methylene disalicylate in feed for animals as follows:

(i) Amount. 70 milligrams per head per day.

(a) Indications for use. Feedlot beef cattle; reduction in the number of liver condemnations due to abscesses.

(b) Limitations. Administer continuously throughout the feeding period.

(i) Amount. 250 milligrams per head per day.

(a) Indications for use. Feedlot beef cattle; reduction in the number of liver condemnations due to abscesses.

(b) Limitations. Administer continuously for 5 days then discontinue for

subsequent 25 days, repeat the pattern during the feeding period.

(3) Bacitracin methylene disalicylate may also be used with:

(i) Amprolium as in § 558.55.

(ii) Amprolium and ethopabate as in § 558.58.

(iii) Carbarosone (not USP) as in § 558.120.

(iv) Decoquinatone alone and with roxarsone as in § 558.195.

(v) Diclazuril alone and with roxarsone as in § 558.198.

(vi) Fenbendazole as in § 558.258.

(vii) Halofuginone hydrobromide alone and with roxarsone as in § 558.265.

(viii) Hygromycin B as in § 588.274.

(ix) Ivermectin as in § 558.300.

§ 558.78

21 CFR Ch. I (4-1-03 Edition)

- (x) Lasalocid sodium alone and with roxarsone as in § 558.311.
- (xi) Monensin alone and with roxarsone as in § 558.355.
- (xii) Narasin alone and with roxarsone as in § 558.363.
- (xiii) Nicarbazine alone and with narasin or narasin and roxarsone as in § 558.366.
- (xiv) Nitarsone as in § 558.369.
- (xv) Robenidine alone and with roxarsone as in § 558.515.
- (xvi) Salinomycin alone and with roxarsone as in § 558.550.
- (xvii) Semduramicin alone and with roxarsone as in § 558.555.
- (xviii) Zoalene alone and with arsanilic acid as in § 558.680.

[41 FR 10993, Mar. 15, 1976]

EDITORIAL NOTE: For FEDERAL REGISTER citations affecting § 558.76, see the List of CFR

Sections Affected, which appears in the Finding Aids section of the printed volume and on GPO Access.

§ 558.78 Bacitracin zinc.

(a) *Approvals.* To sponsors listed in § 510.600(c) of this chapter for use as in paragraph (d) of this section as follows:

(1) No. 046573: 10, 25, 40, and 50 grams per pound as in paragraph (d) of this section.

(2) No. 053389: 5 and 50 grams per pound for chickens as in paragraph (d)(1)(i) of this section.

(b) *Special considerations.* The quantities of antibiotics are expressed in terms of the equivalent amount of antibiotic standard.

(c) *Related tolerances.* See § 556.70 of this chapter.

(d) *Conditions of use.* (1) It is used as follows:

Bacitracin zinc in grams per ton	Combinations in grams per ton	Indications for use	Limitations	Sponsor
(i) 4 to 50		Chickens, turkeys, and pheasants; for increased rate of weight gain and improved feed efficiency.	Growing chickens, turkeys, and pheasants.	046573
(ii) 5 to 20		Quail; for increased rate of weight gain and improved feed efficiency.	Growing quail, feed as the Type C feed to starting quail through 5 weeks of age.	046573
(iii) 10 to 25		Laying chickens; improved feed efficiency and increased egg production		046573
(iv) 10 to 50		Swine; increased rate of weight gain and improved feed efficiency	Growing and finishing swine	046573
(v) 20		Growing-finishing swine; increased rate of weight gain.	In Type C feed	046573
(vi) 20 to 40		Growing-finishing swine; improved feed efficiency.	do	046573

(2) It is used in feed for growing cattle at 35 to 70 milligrams per head per day as follows:

(i) To aid in stimulating growth and improving feed efficiency.

(ii) For increased rate of weight gain and improved feed efficiency; see sponsor 046573.

(3) It may be used as approved in combination with:

(i) Amprolium alone and with roxarsone as in § 558.55.

(ii) Amprolium and ethopabate alone and with roxarsone as in § 558.58.

(iii) Carbarsone as in § 558.120.

(iv) Clopidol alone and with roxarsone as in § 558.175.

(v) Decoquate alone and with roxarsone as in § 558.195.

(vi) Hygromycin B alone and with penicillin as in § 558.274.

(vii) Lasalocid sodium alone or with roxarsone as in § 558.311.

(viii) Monensin alone and with roxarsone as in § 558.355.

(ix) Naracin as in § 558.363.

(x) Robenidine as in § 558.515.

(xi) Salinomycin alone and with roxarsone as in § 558.550.

(xii) Zoalene alone and with arsanilic acid or roxarsone as in § 558.680.

[41 FR 10994, Mar. 15, 1976]

EDITORIAL NOTE: For FEDERAL REGISTER citations affecting § 558.78, see the List of CFR Sections Affected, which appears in the