

**REFERENCES**

Baumgartner M, Brown CA, Secretin MC, van't Hof M, Haschke F: Controlled trials investigating the use of one partially hydrolyzed whey formula for dietary prevention of atopic manifestations until 60 months of age: An overview using meta-analytical techniques. *Nutr Res* 1998;18:1425-1442.

Becker A, Watson W, Ferguson A, Dimich-Ward H, Chan-Yeung M: The Canadian asthma primary prevention study: outcomes at 2 years of age. *J Allergy Clin Immunol* 2004;113:650-656.

Canadian Broadcasting Corporation. The National. The secret life of Dr Chandra. 1/30/06; 1/31/06; 2/1/06. [www.cbc.ca/national/news/chandra](http://www.cbc.ca/national/news/chandra).

Chan YH, Shek LPC, Aw M, Quak SH, Lee BW: Use of hypoallergenic formula in the prevention of atopic disease among Asian children. *J Paediatr Child Health* 2002;38:84-88.

Chan-Yeung M, Manfreda J, Dimich-Ward H, Ferguson A, Watson W, Becker A: A randomized controlled study on the effectiveness of a multifaceted intervention program in the primary prevention of asthma in high-risk infants. *Arch Pediatr Adolesc Med* 2000;154:657-663.

Chandra RK: Five-year follow-up of high-risk infants with family history of allergy who were exclusively breast-fed or fed partial whey hydrolysate, soy and conventional cow's milk formulas. *J Pediatr Gastroenterol Nutr* 1997;24:380-388.

Chandra RK, Hamed A. Cumulative incidence of atopic disorders in high risk infants fed whey hydrolysate, soy, and conventional cow milk formulas. *Ann Allergy* 1991;67:129-132.

Chandra RK, Singh G, Shridhara B: Effect of feeding whey hydrolysate, soy and conventional cow milk formulas on incidence of atopic disease in high risk infants. *Ann Allergy* 1989;63:102-106.

de Seta L, Siani P, Cirillo G, Di Gruttola M, Cimaduomo L, Coletta S: La prevenzione delle malattie allergiche con formula H.A.: follow-up a 24 mesi. Primi risultati. [Prevention of allergic disease by an hypoallergenic formula: preliminary results at the 24 months follow-up]. *Pediatr Med Chir* 1994;16:251-254.

Eigenmann PA, Sicherer SH, Borkowski TA, Cohen BD, Sampson HA: Prevalence of IgE-mediated food allergy among children with atopic dermatitis. *Pediatrics* 1998;101:e8.

Exl BM, Deland U, Secretin MC, Preysch U, Wall M, Shmerling DH: Improved general health status in an unselected infant population following an allergen-reduced dietary intervention programme: the ZUFF-STUDY-PROGRAMME. Part II: infant growth and health status to age 6 months. *Eur J Nutr* 2000;39:145-156.

Exl BM, Deland U, Wall M, Preysch U, Secretin MC, Shmerling DH: Zug-Frauenfeld nutritional survey ("ZUFF study"): allergen-reduced nutrition in a normal infant population and its health-related effects: results at the age of six months. *Nutr Res* 1998;18:1443-1462.

Glassi C, De Sario M, Biggeri A, et al: Changes in prevalence of asthma and allergies among children and adolescents in Italy: 1994-2002. *Pediatrics* 2006;117:34-41.

Maziak W, Behrens T, Brasky TM, et al: Are asthma and allergies in children and adolescents increasing? Results from ISAAC phase I and phase III surveys in Munster, Germany. *Allergy* 2003;58:572-579.

Marini A, Agosti M, Motta G, Mosca F: Effects of a dietary and environmental prevention programme on the incidence of allergic symptoms in high atopic risk infants. Three years follow-up. *Acta Paediatr Suppl* 1996;414:1-22.

Muraro A, Dreborg S, Halken S, Host A, Niggemann B, Aalberse R, Arshad SH, Berg AA, Carlsen KH, Duschek K, Eigenmann P, Hill D, Jones C, Mellon M, Oldeus G, Oranje A, Pascual C, Prescott S, Sampson H, Svartengren M, Vandenplas Y, Wahn U, Warner JA, Warner JO, Wickman M, Zeiger RS: Dietary prevention of allergic diseases in infants and small children. Part III: Critical review of published peer-reviewed observational and interventional studies and final recommendations. *Pediatr Allergy Immunol* 2004;15:291-307.

Osborn DA, Sinn J: Formulas containing hydrolysed protein for prevention of allergy and food intolerance in infants. *Cochrane Database Syst Rev* 2003;CD003664

Sampson HA: Update on food allergy. *J Allergy Clin Immunol* 2004;113:805-19.

Schmidt E, Eden-Kohler J, Tonkaboni F, Tolle J: Alimentary allergy prevention in infants with familial increased allergic risk: the effect of different feeding regimens in the first 6 months on atopic manifestations during the first year of life. A large-scale feeding trial. Nestle Nutrition Workshop Series, Intestinal Immunology and Food Allergy, Raven Press 1995;34:231-248.

Schoetzau A, Gehring U, Wichmann HE: Prospective cohort studies using hydrolysed formulas for allergy prevention in atopy-prone newborns: a systematic review. *Eur J Pediatr* 2001;160:323-332.

Tsai Y-T, Chou C-C, Hsieh K-H: The effect of hypoallergenic formula on the occurrence of allergic diseases in high risk infants. *Acta Paed Sin* 1991;32:137-144.

Vandenplas Y: Atopy at 3 years in high-risk infants fed whey hydrolysate or conventional formula. *Lancet* 1992;339:1118 [letter to editor].

Vandenplas Y, Hauser B, Van den BC, Clybouw C, Mahler T, Hachimi-Idrissi S, Deraeve L, Malfroot A, Dab I: The long-term effect of a partial whey hydrolysate formula on the prophylaxis of atopic disease. *Eur J Pediatr* 1995;154:488-494.

von Berg A, Koletzko S, Grubl A, Filipiak-Pittroff B, Wichmann HE, Bauer CP, Reinhardt D, Berdel D: The effect of hydrolyzed cow's milk formula for allergy prevention in the first year of life: the German Infant Nutritional Intervention Study, a randomized double-blind trial. *J Allergy Clin Immunol* 2003;111:533-540.

Willems R, Duchateau J, Magrez P, Denis R, Casimir G: Influence of hypoallergenic milk formula on the incidence of early allergic manifestations in infants predisposed to atopic diseases. *Ann Allergy* 1993;71:147-150.

Zeiger RS: Prevention of food allergy and atopic disease. *J R Soc Med* 1997;90:21-23.

Zeiger RS, Heller S, Mellon MH, Halsey JF, Hamburger RN, Sampson HA: Genetic and environmental factors affecting the development of atopy through age 4 in children of atopic parents: a prospective randomized study of food allergen avoidance. *Pediatr Allergy Immunol* 1992;3:110-127.