Comments of the Cosmetic, Toiletry, and Fragrance Association (CTFA) Regarding the Scientific and Legal Issues Associated with Nanotechnology in Personal Care Products:

Literature References

Submitted in Regard to Docket Number 2006P-0210: Petition to FDA to Amend its Regulations for Products Composed of Engineered Nanoparticles Generally and Sunscreen Drug Products Composed of Engineered Nanoparticles Specifically

CTFA Comments on Petition to FDA to Amend its Regulations for Products Composed of Engineered Nanoparticles Generally and Sunscreen Drug Products Composed of Engineered Nanoparticles Specifically
Literature References

Barceloux, D. G. (1999), Zinc. *Clinical Toxicology,* 37(2) 279-292.

Beckett, W.S, Chalupa, D.F., Pauly-Brown, A. *et al.* (2005). Comparing inhaled ultrafine versus fine zinc oxide particles in healthy adults; A Human Inhalation Study.*Am. J Respir Crit Care Med.* Vol. 171, 1129-1135.

Bennat, C., and Müller-Goymann, C.C., (2000). Skin penetration and stablization of formulations containing microfine titanium dioxide as physical UV filter. *Int. J. Cosmet. Sci.* **22,** 271-283.

Bermudez, E., Mangum, J.B., Asgharian, A. *et al.* (2002). Long-term pulmonary responses of three laboratory rodent species to subchronic inhalation of pigmentary titanium dioxide particles. *Toxicological Sciences* **70:**86-97.

Bestak, R., and Halliday, G.M., (1996). Sunscreens protect from UV-promoted squambous cell carcinoma in mice chronically irradiated with doses of UV radiation insufficent to cause edema. *J. Photochem, Photobiol. 64(1): 188-193.*

Brendler-Schwaab, S., Czich, A., Epe, B. *et al.* (2004). Photochemical genotoxicity: principles and test methods. Report of a GUM Task Force. *Mutation Research* **566**: 65-91.

Brown, J.S., Zeman, K.L., and Bennett, W.D. (2002). Ultrafine particle deposition and clearance in the healthy and obstructed lung. *Am. J. Respir Crit. Care Med.* **166,**: 1240-1247.

Chen, Y., Chen, J., Dong, J. and Jin, Y. (2004). Comparing study of the effect of nanosized silicon dioxide and microsized silicon dioxide on fibrogenesis in rats. *Toxicol. Ind. Health,* **20:** 21-27.

Dreher, K.L. (2004). Health and environmental impact of nanotechnology: Toxicological assessment of manufactured nanoparticles. *Toxicological Sciences,* **77:** 3-5.

Donaldson, K., Brown, D. Clouter, A *et al.* (2002). The pulmonary toxicology of ultrafine particles. *J. Aerosol Med.* **15:** 213-220.

Dufour, Ek, Kumaravel, T, Nihynek, G.J. *et al.* (2006). Clastogenicity, photo-clastogenicity or pseudo-photo-clastogenicity: Genotoxic effects of zinc oxide in the dark, in pre-irradiated or simultaneously irradiated Chinese hamster ovary cells. *Mutation Research,* Article in Press.

Dunford, R., Salinaro, A, Cai, L. *et al.* (1997). Chemical oxidation and DNA damage catalysed by inorganic sunscreen ingredients. *FEBS Letters,* **418:** 87-90.

Elder, A., Gelein, R., Silva, V., (2006) Translocation of inhaled ultrafine manganese oxide particles to the central nervous system. *Environmental Health Perspectives,* **114**(8); 1172 -1178.

Ferin, J., Oberdörster (1992). Polymer degradation and ultrafine particles: potential inhalation hazards for astronauts. *Acta Astronautical,* **27**: 257-259.

Gamer, A.O., Leibold, E., vanRavenzwaay B. (2006). The in vitro absorption of microfine zinc oxide and titanium dioxide through porcine skin. *Toxicology in Vitro,* **20**: 301-307.

Gasparro, F., Mitchnick, M., and Nash, J.F. (1998). A review of sunscreen safety and efficacy. *J. Photochem. Photobiol.* **68**(3): 243-356.

Geiser, M., Rothen-Rutishauser, B., Kapp, N. *et al.* (2005). Ultrafine particles cross cellular membranes by nonphagocytic mechanisms in lungs and cultured cells. *Environmental Health Perspectives,* **113**(11): 1555-1560.

Gopee, N.V., Roberts, D. W., Webb, P, Cozart, C., *et al.* (2006). Penetration of Nanoscale quantum dots in dermabraded mouse skin. *2006 FDA Science Forum, Poster Abstract,* **B-36,** March 2006. At http://www.cfsan.fda.gov

Gurr, J-R., Wang, A.S, Chen- Hung, C, *et al.* (2005). Ultrafine titanium dioxide particles in the absence of photoactivation can induce oxidative damage to human bronchial epithelial cells. *Toxicology* **213**: 66-73.

Hardman, R., (2006). A toxicologic review of quantum dots: toxicity depends on physicochemical and environmental factors. *Environmental Health Perspectives,* **114**(2): 165-172.

Höhr, D., Steinfartz,Y., Schins, R., Knaapen, Ad. M., *et al.* (2002) The surface area rather than the surface coating determines the acute inflammatory response after instillation of fine and ultrafine TiO_2 in the rat. *Int. J. Hyg. Environ. Health,* **205**: 239-244.

Hostynek, J.J., Factors determining percutaneous metal absorption, (2003). *Food and Chemical Toxicology,* **41** 327-345.

Janssen, Y.M., Marsh, J.P., Driscoll, K.E., *et al.* (1994). Increased expression of manganese-containing superoxide dismutase in rat lung after inhalation of inflammatory and fibrogenic materials. *Free Radical Biology & Medicine,* **16**(3): 315-322.

Lademann, J.M Weigmann, H.J., Rickmeyer, C., *et al.* (1999). Penetration of Titanium Dioxide Microparticles in a sunscreen formulation into the horny layer and follicular orifice. *Skin Pharmacol Appl Skin Physiol*, **12**:247-256.

Menzel, F., Reiner, T., Vogt, J., Butz, T. (2004). Investigations of percutaneous uptake of ultrafine TiO_2 particles at the high energy ion nanoprobe LIPSION. *Nuclear Instruments and Methods in Phyysics*, **B** 219-220, 82-86.

Mills, A., LeHunte, S., (1997). An overview of semiconductor photocatalysis. *J. Photochem Photobiol.,* **108**: 1-35.

Mills, N.L., Robinson, A.M., Davies, A.D., *et al.* (2006). Do inhaled carbon nanoparticles translocate directly into the circulation in humans? *Am. J. Respir Crit Care Med* **173**(4):426-431.

Moloney, F.J., Collins, S., Murphy, G.M., (2002). Sunscreens: safety, efficacy and appropriate use. *Am. J. Clin. Dermatol.* **3** (3): 185-191.

Nakagawa, Y., Wakuri, S., Sakamoto, K. Tanaka, N., (1997). The photogenotoxicity of titanium dioxide particles. *Mutation Reserach,* **394**: 125-132.

Nash, J.F., (2006). Human Safety and Efficacy of Ultraviolet Filters and Sunscreen Products. *Dermatol. Clin.* **24**: 35-51.

Nel, A., Xia, T., Madler, L., Ning, L. (2006). Toxic potential of materials at the nanolevel. *Science*, **311**;622-627.

Nemar, A., Vanbilloen, H., Hoylaerts, M.F. *et al.* (2001). Passage of inhaled particles into the blood circulation in humans. *Circulation* **105:** 411-414.

Nohynek, G.J., Schefer, H., (2001) Benefits and risk of organic ultraviolet filters. *Regulatory Toxicology and Pharmacology,* **33**: 285-299.

Oberdörster, G., Ferin, J., Gelein, R. *et al.* (1992). Role of the alveolar macrophage in lung injury: studies with ultrafine particles. *Environmental Health Perspectives*, **97**: 193-199.

Oberdörster, G., (1996). Significance of particle parameters in the evaluation of exposure-dose-response relationships of inhaled particles. *Inhalation Toxicology* **8**(suppl): 73-89.

Oberdörster, G., Oberdörster, E., Oberdörster, J. (2005). Nanotoxicology: An emerging discipline evolving from studies of ultrafine particles. *Environmental Health Perspectives*, **1131**(7): 823-839.

Pflücker, F., Hohenberg, H., Hölzle, E., *et al.* (1999). The outermost stratum corneum layer is an effective barrier against dermal uptake of topically applied micronized titanium dioxide. *Intl. J. Cosmet. Sci.* **21**:399-411.

Pflücker, F., Wendel, V., Hohenberg, H. *et al.* (2001). The human stratum corneum layer: An effective barrier against dermal uptake of different forms of topically applied micronized titanium dioxide. *Skin Pharmacol Appl Skin Physiol* **14**(suppl 1): 92-97.

Pinnell, S.R., Fairhurst, D., Gillies, R., *et al.* (2000). Microfine zinc oxide is a superior sunscreen ingredient to microfine titanium dioxide. *Dermatol. Surg.* **26:**309-314.

Renwick, L.C., Brown, D., Clouter, A., Donaldson, K., (2004). Increased inflammation and altered macrophage chemotactic responses caused by two ultrafine particle types. *Occup. Environ. Med.* **61:**442-447.

Ryman-Rasmussen, J.P., Riciere, J.E., Monteiro-Riviere, N.A., (2006). Penetration of intact skin by quantum dots with diverse physicochemical properties. *Toxicological Sciences* **91**(1): 159-165.

Sayes, C.M., Wahi, R., Kurian, F.A., *et al.* (2006). Correlating nanoscale tiania structure with toxicity: A cytotoxicity and inflammatory response study with human dermal fibroblasts and human lung epithelial cells. *Toxicologial Sciences* **92**(1): 174-185.

Schulz, J., Hohenberg, H., Pflücker, *et al.* (2002). Distribution of sunscreens on skin. *Advanced Drug Delivery Review* **54** (suppl 1): S157-S163.

Semmler, M., Seitz, J., Erbe, F., *et al.* (2004). Long-term clearance kinetics of inhaled ultrafine insoluble iridium particles from the rat lung, including transient translocation into secondary organs. *Inhal. Toxicol.* **16**(6-7):453-459.

Suzuki, M., (1987). Protective effect of fine-particle titanium dioxide on UVB-induced DNA damage in hairlesss mouse skin. *Photodermatology* **4**:209-211.

Tan, M.-H., Commens, C.A., Burnett, L., Snitch, P.J., (1996). A pilot study of the percutaneous absorption of microfine titanium dioxide from sunscreens, *Aust. J. Dermatol.* **37**:185-187.

Tinkle, S.S., Antonini, J.M., Rich, B.A., *et al.* (2003). Skin as a route of exposure and sensitization in chronic beryllium disease. *Environmental Health Perspective* **111:**1202-1208.

Van der Molen, R.G., Hurks, H.M.H., Out-Luiting, C., *et al.* (1998). Efficacy of micronized titanium dioxide-containing compounds in protection against UVB-induced immunosuppression in humans in vivo. *J. Photochem. Photobiol.* **44**:(143-150).

Walter, J.F., Dequoy, P.R., (1980) The hairless mouse as a model for evaluating sunscreens. *Arch Dermatol* **116**:419 - 421.

Warheit, D.B., Webb, T.R., Sayes, C.M. *et al.* (2006). Pulmonary instillation studies with nanoscale TiO_2 rods and dots in rats: Toxicity is not dependent upon particle size and surface area. *Toxicologial Sciences* **91**(1):227-236.

Wiebert, P., Sanchez-Crespo, A., Falk, R. *et al.* (2006). Negligble clearance of ultrafine particles retained in healthy and affected human lungs. *Eur Respir J, 28:286-290.*

Wiebert, P., Sanchez-Crespo, A., Falk, R. *et al.* (2006). No significant translocation of inhaled 35-nm carbon particles to the circulation in humans. *Inhal Toxicol.* **18**(10): 741-747.

Yamamoto, A., Honma, R., Sumita, M., Hanawa, T., (2003). Cytotoxicity evaluation of ceramic particles of different sizes and shapes. *J. Biomed. Mater. Res.***68**: 244-256.