



# ALCLOMETASONE DIPROPIONATE LOTION 0.05%

R<sub>x</sub> only


**For Dermatologic Use Only—  
Not for Ophthalmic Use.**

## DESCRIPTION

Alclometasone Dipropionate Lotion, 0.05% contains alclometasone dipropionate (7 $\alpha$ -chloro-11 $\beta$ ,17,21-trihydroxy-16 $\alpha$ -methylpregna-1,4-diene-3,20-dione 17,21-dipropionate), a synthetic corticosteroid for topical dermatologic use. The corticosteroids constitute a class of primarily synthetic steroids used topically as anti-inflammatory and antipruritic agents.

Chemically, alclometasone dipropionate is C<sub>28</sub>H<sub>37</sub>ClO<sub>7</sub>. It has the following structural formula:

Alclometasone dipropionate has the molecular weight of 521. It is a white powder,



insoluble in water, slightly soluble in propylene glycol, and moderately soluble in hexylene glycol.

Each ml contains 0.5 mg of alclometasone dipropionate in a lotion consisting of stearyl alcohol, cetyl alcohol, glyceryl monostearate, sorbitan monooleate, mineral oil, purified water, monobasic sodium phosphate, phosphoric acid, polysorbate 20, simethicone and propylene glycol.

## CLINICAL PHARMACOLOGY

Like other topical corticosteroids, alclometasone dipropionate has anti-inflammatory, antipruritic, and vasoconstrictive properties. The mechanism of the anti-inflammatory activity of the topical steroids, in general, is unclear. However, corticosteroids are thought to act by the induction of phospholipase A<sub>2</sub> inhibitory proteins, collectively called lipocortins. It is postulated that these proteins control the biosynthesis of potent mediators of inflammation such as prostaglandins and leukotrienes by inhibiting the release of their common precursor, arachidonic acid. Arachidonic acid is released from membrane phospholipids by phospholipase A<sub>2</sub>.

**Pharmacokinetics:** The extent of percutaneous absorption of topical corticosteroids is determined by many factors, including the vehicle and the integrity of the epidermal barrier. Occlusive dressings with hydrocortisone for up to 24 hours have not been demonstrated to increase penetration; however, occlusion of hydrocortisone for 96 hours markedly enhances penetration. Topical corticosteroids can be absorbed from normal intact skin. Inflammation and/or other disease processes in the skin may increase percutaneous absorption. A study utilizing a radiolabeled alclometasone dipropionate ointment formulation was performed to measure systemic absorption and excretion. Results indicated that approximately 3% of the steroid was absorbed during 8 hours of contact with intact skin of normal volunteers.

Studies performed with alclometasone dipropionate indicate that these products are in the low to medium range of potency as compared with other topical corticosteroids.

## INDICATIONS AND USAGE

Alclometasone Dipropionate Lotion, 0.05% is a low to medium potency corticosteroid indicated for the relief of the inflammatory and pruritic manifestations of corticosteroid-responsive dermatoses. Alclometasone Dipropionate Lotion, 0.05% may be used in pediatric patients 1 year of age or older, although the safety and efficacy of drug use for longer than 3 weeks have not been established (see PRECAUTIONS: Pediatric Use). Since the safety and efficacy of Alclometasone

Dipropionate Lotion, 0.05% has not been established in pediatric patients below 1 year of age, their use in this age-group is not recommended.

## CONTRAINDICATIONS

Alclometasone Dipropionate Lotion, 0.05% is contraindicated in those patients with a history of hypersensitivity to any of the components in these preparations.

## PRECAUTIONS

**General:** Systemic absorption of topical corticosteroids can produce reversible hypothalamic-pituitary-adrenal (HPA) axis suppression with the potential for glucocorticosteroid insufficiency after withdrawal of treatment. Manifestations of Cushing syndrome, hyperglycemia, and glucosuria can also be produced in some patients by systemic absorption of topical corticosteroids while on treatment. Patients applying a topical steroid to a large surface area or to areas under occlusion should be evaluated periodically for evidence of HPA axis suppression. This may be done by using the ACTH stimulation, A.M. plasma cortisol, and urinary free cortisol tests.

The effects of alclometasone dipropionate on the HPA axis have been evaluated. In one study, alclometasone dipropionate was applied to 30% of the body twice daily for 7 days, and occlusive dressings were used in selected patients either 12 hours or 24 hours daily. In another study, alclometasone dipropionate was applied to 80% of the body surface of normal subjects twice daily for 21 days with daily 12-hour periods of whole body occlusion. Average plasma and urinary free cortisol levels and urinary levels of 17-hydroxysteroids were decreased (about 10%), suggesting suppression of the HPA axis under these conditions. Plasma cortisol levels have also been demonstrated to decrease in pediatric patients treated twice daily for 3 weeks without occlusion.

If HPA axis suppression is noted, an attempt should be made to withdraw the drug, to reduce the frequency of application, or to substitute a less potent corticosteroid. Recovery of HPA axis function is generally prompt upon discontinuation of topical corticosteroids. Infrequently, signs and symptoms of glucocorticosteroid insufficiency may occur, requiring supplemental systemic corticosteroids. For information on systemic supplementation, see prescribing information for those products.

Pediatric patients may be more susceptible to systemic toxicity from equivalent doses due to their larger skin surface area to body mass ratios (see PRECAUTIONS: Pediatric Use).

If irritation develops, Alclometasone Dipropionate Lotion, 0.05% should be discontinued and appropriate therapy instituted. Allergic contact dermatitis with corticosteroids is usually diagnosed by observing a *failure to heal* rather than noting a clinical exacerbation, as with most topical products not containing corticosteroids. Such an observation should be corroborated with appropriate diagnostic patch testing.

If concomitant skin infections are present or develop, an appropriate antifungal or antibacterial agent should be used. If a favorable response does not occur promptly, use of Alclometasone Dipropionate Lotion, 0.05% should be discontinued until the infection has been adequately controlled.

**Information for Patients:** Patients using topical corticosteroids should receive the following information and instructions:

1. This medication is to be used as directed by the physician. It is for external use only. Avoid contact with the eyes.
2. This medication should not be used for any disorder other than that for which it was prescribed.
3. The treated skin area should not be bandaged, otherwise covered or wrapped so as to be occlusive, unless directed by the physician.
4. Patients should report to their physician any signs of local adverse reactions.
5. Parents of pediatric patients should be advised not to use Alclometasone Dipropionate Lotion, 0.05% in the treatment of diaper dermatitis. Alclometasone Dipropionate Lotion, 0.05% should not be applied in the diaper

## Alclometasone Dipropionate Lotion, 0.05%

area as diapers or plastic pants may constitute occlusive dressing (see DOSAGE AND ADMINISTRATION).

- This medication should not be used on the face, underarms, or groin areas unless directed by the physician.
- As with other corticosteroids, therapy should be discontinued when control is achieved. If no improvement is seen within 2 weeks, contact the physician.

**Laboratory Tests:** The following tests may be helpful in evaluating patients for HPA axis suppression:  
ACTH stimulation test  
A.M. plasma cortisol test  
Urinary free cortisol test

**Carcinogenesis, Mutagenesis, Impairment of Fertility:** Long-term animal studies have not been performed to evaluate the carcinogenic potential or the effect on fertility of topical corticosteroids.

**Pregnancy: Teratogenic Effects:** Pregnancy Category C. Corticosteroids have been shown to be teratogenic in laboratory animals when administered systemically at relatively low dosage levels. Some corticosteroids have been shown to be teratogenic after dermal application in laboratory animals. There are no adequate and well-controlled studies in pregnant women. Alclometasone Dipropionate Lotion, 0.05% should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

**Nursing Mothers:** Systemically administered corticosteroids appear in human milk and could suppress growth, interfere with endogenous corticosteroid production, or cause other untoward effects. It is not known whether topical administration of topical corticosteroids could result in sufficient systemic absorption to produce detectable quantities in human milk. Because many drugs are excreted in human milk, caution should be exercised when Alclometasone Dipropionate Lotion, 0.05% is administered to a nursing woman.

**Pediatric Use:** Alclometasone Dipropionate Lotion, 0.05% may be used with caution in pediatric patients 1 year of age or older, although the safety and efficacy of drug use for longer than 3 weeks have not been established. Use of alclometasone dipropionate is supported by results from adequate and well-controlled studies in pediatric patients with corticosteroid-responsive dermatoses. Since the safety and efficacy of Alclometasone Dipropionate Lotion, 0.05% has not been established in pediatric patients below 1 year of age, its use in this age-group is not recommended. Because of a higher ratio of skin surface area to body mass, pediatric patients are at a greater risk than adults of HPA axis suppression and Cushing syndrome when they are treated with topical corticosteroids. They are therefore also at greater risk of adrenal insufficiency during and/or after withdrawal of treatment. Adverse effects, including striae, have been reported with inappropriate use of topical corticosteroids in infants and children. Pediatric patients applying Alclometasone Dipropionate Lotion, 0.05% to >20% of the body surface area are at higher risk for HPA axis suppression. HPA axis suppression, Cushing syndrome, linear growth retardation, delayed weight gain, and intracranial hypertension have been reported in pediatric patients receiving topical corticosteroids. Manifestations of adrenal suppression in pediatric patients include low plasma cortisol levels and absence of response to ACTH stimulation. Manifestations of intracranial hypertension include bulging fontanelles, headaches, and bilateral papilledema.

Alclometasone Dipropionate Lotion, 0.05% should not be used in the treatment of diaper dermatitis.

**Geriatric Use:** A limited number of patients at or above 65 years of age have been treated with alclometasone dipropionate in US clinical trials. The number of patients is too small to permit separate analysis of efficacy and safety. The single adverse reaction reported with alclometasone dipropionate in this population was similar to those reactions reported by younger patients. Based on available data, no adjustment of dosage of Alclometasone Dipropionate Lotion, 0.05% in geriatric patients is warranted.

### ADVERSE REACTIONS

The following local adverse reactions have been reported with Alclometasone Dipropionate Lotion, 0.05% in approximately 2% of patients: itching and burning, erythema, dryness, irritation, and papular rashes.

The following additional local adverse reactions have been reported infrequently with topical corticosteroids, but may occur more frequently with the use of occlusive dressings. These reactions are listed in approximate decreasing order of occurrence: folliculitis, acneiform eruptions, hypopigmentation, perioral dermatitis, allergic contact dermatitis, secondary infection, skin atrophy, striae, and miliaria.

### OVERDOSAGE

Topically applied Alclometasone Dipropionate Lotion, 0.05% can be absorbed in sufficient amounts to produce systemic effects (see PRECAUTIONS).

### DOSAGE AND ADMINISTRATION

Apply a thin film of Alclometasone Dipropionate Lotion, 0.05% to the affected skin areas 2 or 3 times daily; massage gently until the medication disappears. Alclometasone Dipropionate Lotion, 0.05% may be used in pediatric patients 1 year of age or older. Safety and effectiveness of Alclometasone Dipropionate Lotion, 0.05% in pediatric patients for more than 3 weeks of use have not been established. Use in pediatric patients under 1 year of age is not recommended. As with other corticosteroids, therapy should be discontinued when control is achieved. If no improvement is seen within 2 weeks, reassessment of diagnosis may be necessary.

Alclometasone Dipropionate Lotion, 0.05% should not be used with occlusive dressings unless directed by a physician. Alclometasone Dipropionate Lotion, 0.05% should not be applied in the diaper area if the child still requires diapers or plastic pants as these garments may constitute occlusive dressing.

**Geriatric Use:** In studies where geriatric patients (65 years of age or older, see PRECAUTIONS) have been treated with alclometasone dipropionate, safety did not differ from that in younger patients; therefore, no dosage adjustment is recommended.

### HOW SUPPLIED

Alclometasone Dipropionate Lotion, 0.05% is supplied in 2oz and 4oz bottles. Store between 2°C and 30°C (36°F and 86°F).

R<sub>x</sub> only NDC 1  
59 mL (2 Fl Oz)

**Alclometasone  
Dipropionate  
Lotion 0.05%**

For dermatologic use only  
Not for ophthalmic use

**SHAKE WELL BEFORE USING.**  
Store between 2°C and 30°C  
(36°F and 86°F).  
**WARNING: Keep out of  
reach of children.**

**USUAL DOSAGE:** Apply a thin  
film to the affected areas two or  
three times daily.  
See package insert for complete  
prescribing information.

Each ml contains 0.5 mg of  
alclometasone dipropionate in a  
lotion consisting of stearyl alcohol,  
cetyl alcohol, glyceryl  
monostearate, sorbitan  
monopalmitate, mineral oil,  
purified water, monobasic sodium  
phosphate, phosphoric acid,  
polysorbate 20, simethicone and  
propylene glycol.

This product sealed for your  
protection.  
If the seal is broken or missing  
return to place of purchase.

R<sub>x</sub> only NDC 1  
59 mL (2 Fl Oz)

**Alclometasone  
Dipropionate  
Lotion 0.05%**

For dermatologic use only  
Not for ophthalmic use

**SHAKE WELL BEFORE USING.**  
Store between 2°C and 30°C  
(36°F and 86°F).  
**WARNING: Keep out of  
reach of children.**

**USUAL DOSAGE:** Apply a thin  
film to the affected areas two or  
three times daily.  
See package insert for complete  
prescribing information.

Each ml contains 0.5 mg of  
alclometasone dipropionate in a  
lotion consisting of stearyl alcohol,  
cetyl alcohol, glyceryl  
monostearate, sorbitan  
monopalmitate, mineral oil,  
purified water, monobasic sodium  
phosphate, phosphoric acid,  
polysorbate 20, simethicone and  
propylene glycol.

This product sealed for your  
protection.  
If the seal is broken or missing  
return to place of purchase.

P????  
R7/06



3

R<sub>x</sub> only

NDC  
118 mL (4 Fl Oz)

**Alclometasone  
Dipropionate  
Lotion 0.05%**

For dermatologic use only  
Not for ophthalmic use

**SHAKE WELL BEFORE USING.**  
Store between 2°C and 30°C  
(36°F and 86°F).  
**WARNING: Keep out of  
reach of children.**

**USUAL DOSAGE:** Apply a thin  
film to the affected areas two or  
three times daily.  
See package insert for complete  
prescribing information.

Each ml contains 0.5 mg of  
alclometasone dipropionate in a  
lotion consisting of stearyl alcohol,  
cetyl alcohol, glyceryl  
monostearate, sorbitan  
monopalmitate, mineral oil,  
purified water, monobasic sodium  
phosphate, phosphoric acid,  
polysorbate 20, simethicone and  
propylene glycol.

This product sealed for your  
protection.  
If the seal is broken or missing  
return to place of purchase.

R<sub>x</sub> only

NDC  
118 mL (4 Fl Oz)

**Alclometasone  
Dipropionate  
Lotion 0.05%**

For dermatologic use only  
Not for ophthalmic use

**SHAKE WELL BEFORE USING.**  
Store between 2°C and 30°C  
(36°F and 86°F).  
**WARNING: Keep out of  
reach of children.**

**USUAL DOSAGE:** Apply a thin  
film to the affected areas two or  
three times daily.  
See package insert for complete  
prescribing information.

Each ml contains 0.5 mg of  
alclometasone dipropionate in a  
lotion consisting of stearyl alcohol,  
cetyl alcohol, glyceryl  
monostearate, sorbitan  
monopalmitate, mineral oil,  
purified water, monobasic sodium  
phosphate, phosphoric acid,  
polysorbate 20, simethicone and  
propylene glycol.

This product sealed for your  
protection.  
If the seal is broken or missing  
return to place of purchase.

P????  
R7/06



3

1

100% 3 x .875

**Rx only**  
**Alclometasone  
Dipropionate  
Lotion 0.05%**

**Sample - Not for Sale**  
**SHAKE WELL BEFORE USING.**  
**WARNING: Keep out of reach of children.**  
Store between 2°C and 30°C (36°F and 86°F).  
**USUAL DOSAGE:** Apply a thin film to the affected areas two or three times daily. See insert for complete prescribing information.  
Each ml contains 0.5 mg of alclometasone dipropionate in a lotion consisting of stearyl alcohol, cetyl alcohol, glyceryl monostearate, sorbitan monopalmitate, mineral oil, purified water, monobasic sodium phosphate, phosphoric acid, polysorbate 20, stearic acid and propylene glycol.  
This product sealed for your protection. If the seal is missing or broken return to place of purchase.  
L740503 R7/06

LOT NO

EXP

200%

**Rx only**  
**Alclometasone  
Dipropionate  
Lotion 0.05%**  
For dermatologic use only  
Not for ophthalmic use

**Sample - Not for Sale**  
**SHAKE WELL BEFORE USING.**  
**WARNING: Keep out of reach of children.**  
Store between 2°C and 30°C (36°F and 86°F).  
**USUAL DOSAGE:** Apply a thin film to the affected areas two or three times daily. See insert for complete prescribing information.  
Each ml contains 0.5 mg of alclometasone dipropionate in a lotion consisting of stearyl alcohol, cetyl alcohol, glyceryl monostearate, sorbitan monopalmitate, mineral oil, purified water, monobasic sodium phosphate, phosphoric acid, polysorbate 20, simethicone and propylene glycol.  
This product sealed for your protection. If the seal is missing or broken return to place of purchase.  
L740503 R7/06

LOT NO

EXP

EXP

LOT

R-only

59 mL (2 Fl Oz)

**Alclometasone  
Dipropionate  
Lotion 0.05%**

For dermatologic use only  
Not for ophthalmic use

**SHAKE WELL BEFORE USING.**

Store between 2°C and 30°C  
(36°F and 86°F).

**WARNING:** Keep out of reach of children.  
**USUAL DOSAGE:** Apply a thin film to the  
affected areas two or three times daily. See  
package insert for complete prescribing  
information.

Each mL contains 0.5 mg of alclometasone  
dipropionate in a lotion consisting of stearyl  
alcohol, cetyl alcohol, glyceryl monolearate,  
sorbitan monopalmitate, mineral oil, purified  
water, monobasic sodium phosphate,  
phosphoric acid, polysorbate 20, simethicone  
and propylene glycol.

This product sealed for your protection.  
If the seal is broken or missing return to  
place of purchase.



3

LOT  
EXP

R<sub>x</sub> only

NDC  
118 mL (4 Fl Oz)

**Alclometasone  
Dipropionate  
Lotion 0.05%**

For dermatologic use only

Not for ophthalmic use



**SHAKE WELL BEFORE USING.**  
Store between 2°C and 30°C  
(36°F and 86°F).

**WARNING: Keep out of reach of children.**  
**USUAL DOSAGE:** Apply a thin film to the  
affected areas two or three times daily.  
See package insert for complete  
prescribing information.

Each ml contains 0.5 mg of alclometasone  
dipropionate in a lotion consisting of stearyl  
alcohol, cetyl alcohol, glyceryl monostearate,  
sorbitan monopalmitate, mineral oil, purified  
water, monobasic sodium phosphate,  
phosphoric acid, polysorbate 20, simethicone  
and propylene glycol.

This product sealed for your protection.  
If the seal is broken or missing return to place  
of purchase.

L740504  
R7/06

