TITLE

RESEARCH PAPER (-)-HYDROXYCITRIC ACID - THE PRINCIPAL ACID IN THE FRUITS OF GARCINIA CAMBOGIA DESR.

Y. S. Lewis and S. Neelakantan
Central Food Technological Research Institute, Mysore, India
Received 15 March 1964. Available online 2 April 2001.

Summary

The principal acid in the highly acidic fruits of Garcinia Cambogia has been identified as (−)-hydroxycitric acid. This acid has not been encountered in nature before. Fruits of two other species of Garcinia were found to contain the same acid, and the leaves of Hibiscus cannabinus, (+)-allo-hydroxycitric acid, an isomer reported earlier in Hibiscus sabdariffa. Wide occurrence of Hydroxycitric acid in nature is thus indicated.

CONCLUSION OF STUDY

The study presents evidence that (-)-Hydroxycitric Acid is the principal acid present in the fruit rinds of Garcinia Cambogia. Although four isomers of Hydroxycitric acid are possible, this study confirms that only the (-)-isomer occurs in the fruits of Garcinia Cambogia. The presence of (-)-Hydroxycitric Acid had not been confirmed in nature prior to the publication of this article.

HOW THIS STUDY IS RELEVANT TO OUR PRODUCT

The study describes the specific isomer of Hydroxycitric acid that is present in the fruit rind of Garcinia Cambogia as (-)-Hydroxycitric acid. This particular ingredient is widely used in India for culinary purposes. The (-)-Hydroxycitric acid isomer is identified in the study is the functional ingredient in our Trim products.

We deliver this as a stable Potassium Salt of (-)-Hydroxycitric Acid. Various salts such as a Potassium Salt, a Potassium / Calcium Salt, a Calcium Salt etc are commonly sold in the USA as ingredients in dietary supplements and mainstream beverages.

RESEARCH PAPER (-)-HYDROXYCITRIC ACID - THE PRINCIPAL ACID IN THE FRUITS OF GARCINIA CAMBOGIA DESR.

Y. S. Lewis and S. Neelakantan
Central Food Technological Research Institute, Mysore, India
Received 15 March 1964. Available online 2 April 2001.

Abstract

The principal acid in the highly acidic fruits of Garcinia Cambogia has been identified as (−)-hydroxycitric acid. This acid has not been encountered in nature before. Fruits of two other species of Garcinia were found to contain the same acid, and the leaves of Hibiscus cannabinus, (+)-allo-hydroxycitric acid, an isomer reported earlier in Hibiscus sabdariffa. Wide occurrence of Hydroxycitric acid in nature is thus indicated.

Introduction

Very little information is available in literature about the Hydroxycitric acids (four isomers are possible, since two asymmetric C atoms are present). Lippmann reported the presence of an acid in beet juice which he suspected to be Hydroxycitric acid; but the acid was optically inactive. Pawolleck had synthesized a salt of the Hydroxycitric acid earlier, and more recently Martius succeeded in preparing all the four isomers, starting from trans-aconitic acid.

Griebe isolated a Hydroxycitric acid from *Hibiscus Sabdariffa* Linn. calyxes and identified it as (+) allo-hydroxycitric acid lactone, described by Martius The dried rind of the fruit of Garcinia Cambogia Desr., popularly known as "*Malabar Tamarind*", is extensively used all over the west coast of South India for culinary purposes, and commercially for "*Colombo curing*" of fish. The fruit contains about 30 per cent acid calculated as citric acid on dry wt. basis. The organic acids present have been mistakenly identified in the past as tartaric and citric acids. The evidence which has led to the identification of the acid as (-) Hydroxycitric acid is presented here. This acid has so far not been encountered in nature.

Experimental And Results

Both the dried rind of the fruit, as available commercially, and fresh fruits were used for the investigations. The fresh succulent fruit is about the size of an orange, with a thin skin and deep vertical furrows forming blunt lobes.

Paper Chromatography Using extracts of the fruit, two predominant acid spots were found on chromatograms run in different solvent systems. The Rf values of unknowns were mostly so close to tartaric and citric acids (Table 1) that they were apt to be mistaken for them. A spray of 5% sodium metavanadate coloured the lower acid spot reddish orange: this reaction is supposed to be a characteristic of optically active tartic acid. However, attempts at quantitative determination of tartaric and citric acids by A.O.A.C. methods failed to give any potassium bitartrate or pentabromacetone precipitates, even when large aliquots were taken.

TABLE 1. R_f OF ACID SPOTS

Spot	R_f in			
	BAW	BFW	EFW	PAW
No. 1 (free acid)	0·34	0·25	0·22	0·36
Tartaric	0·33	0·24	0·34	0·27
No. 2 (lactone)	0·46	0·42	0·49	0·26
Citric	0·45	0·40	0·48	0·18

BAW—*n*-Butanol: acetic: water (4:1:5), Whatman No. 1 paper.
BFW—*n*-Butanol: formic: water (4:1:5), Whatman No. 2 paper.
EFW—Ether: formic acid: water (10:1:5), Whatman 3MM paper.
PAW—Propanol: ammonia: water (6:3:1), Whatman No. 2 paper.

Column Chromatography

Gradient elution of acids in the fruit, juice, using a Dowex formate column gave a pattern (Fig. I) showing the largest peak in the position occupied by citric and isocitric acids (also malonic and pyruvic). However, the pooled fractions analysed for citric acid by the isocitric dehydrogenase method showed complete absence of this acid. The other acids were ruled out by a consideration of the R_f, values on chromatograms.

Isolation of Acid

Attempts to separate the two acids by fractional precipitation of the calcium or barium salts proved futile. Separation by quantitative paper chromatography by application of a concentrated band at the starting line gave two separate bands. But when these bands were eluted, concentrated and tested again, each gave the original two spots. Ether extraction of the mixture of acids showed a partial concentration of the upper spot, giving a clue that lactone might be involved. On saponification of the mixture using excess alkali and passing through a cation-exchange resin (Zeocarb 2 15) the eluate showed only the lower spot. But on concentration over a water bath, the upper spot also appeared. On heating the concentrate over a water bath for several hours and keeping in a desiccator for several hours a crude crystalline material was obtained. This gave only one spot (upper) on chromatograms. On titration of this material with alkali, two different end-points were obtained, using phenolphthaltin. In the cold and after boiling, characteristic of lactones. It was thus clear that the two spots on chromatograms are those of a y-hydroxy acid and its lactone.

(-)-Hydroxycitric acid 621



FIG. 1. DIAGRAM OF THE ION-EXCHANGE SEPARATION OF THE ORGANIC ACIDS OF *Garcinia cambogia* ON DOWEX 1 FORMATE COLUMN.

Isolation of the acid on a large scale was carried out by the following procedure. The dried rind was cooked with about three volumes of water in an autoclave (10 lb sq in for 15 min) and the extract filtered first through cloth and then paper. The filtrate was concentrated to a small volume and treated with two volumes of alcohol to remove pectin. The clear filtrate was neutralized with alkali, when a viscous, dark, heavy liquid separated (it had been found by previous experience that this oily liquid is the highly hygroscopic alkali salt of the acid). The clear supernatant was decanted off and the oily liquid washed with 60 % alcohol several times.

By repeated treatment with absolute alcohol, this could be dried to a pale yellow hygroscopic powder, which formed the pure alkali salt. Aqueous solutions of the alkali salt were passed

through a cation-exchange resin (Zeocarb 215) for recovery of the acid. The acid itself could not be crystallized because evaporation resulted in formation of the lactone. The syrup obtained on evaporation of the eluate from the ion-exchange resin was dried on a water bath for several hours, and then in a vacuum oven at 100" for g-10 hr. A crude crystalline mass was obtained after keeping in a desiccator for several days. Alternately, the material obtained by evaporation on a water bath could be seeded with a few crystals of the lactone. The crude crystalline material, on extraction and re-crystallization from ether, gave clusters of small needle-shaped crystals of the lactone (Fig. 2).

Characteristics of the Acid

Some of the characteristics of the lactone and the acid are shown in Table 2. The rotation for acid was obtained by saponifying the lactone with excess alkali and adding an equivalent amount of HCl before taking the reading. The equivalent weight of pure lactone is 69, as determined by alkali titration or silver salt decomposition.
By titration and measurement of conductivity the acid proved to be a 3c_hydroxy tribasic acid.

FIG. 2. CRYSTALS OF HYDROXYCITRIC ACID LACTONE.

TABLE 2. COMPARISON OF "GARCINIA" ACID WITH "HIBISCUS" ACID

Property	Hibiscus acid (lactone)	Garcinia acid (lactone)
m.p.	183°	178°
$[\alpha]_D^{20}$	−122	−100
For free acid	+ 31	− 20
After saturation with Borax	+ 31	− 92
Sublimation	−	−
Crystal shape	Needles	Needles
Hygroscopicity	High	Slight
Solubility	High in water and alcohol; slight in ether	High in alcohol and water, fair in ether
Action of dehydrogenase (cucumber seed or liver homogenate) on free acid	Negative	Negative
Paper chromatography R_fs in:		
BFW* — Acid	0·15	0·24
Lactone	0·39	0·42
PAW — Acid	0·35	0·36
Lactone	0·26	0·26
Metavanadate spray		
Acid spot	Yellow	Reddish orange
Lactone spot	Yellow	Yellow
Quinine salt, m.p.	227° decomp.	215° decomp.

* See Table 1.

The molecular weight of lactone from alkali titres is 190; of the acid, 208.

The i.r. spectra of the ethyl ester (b.p. = 178° at 2 mm), showed ester groups (5·41–5·76 μ), OH groups (2·74–2·79 μ) and a γ-lactone (5·62–5·68 μ). The ester analysed for $C_{10}H_{14}O_7$ (Found: C, 49·2; H, 6·0. $C_{10}H_{14}O_7$ required: C, 48·78; H, 5·73%) and had a saponification value of 686. The lactone gave analysis (Found: C, 38·2; H, 3·4. $C_8H_6O_7$ required: C, 37·9; H, 3·15%) for $C_8H_6O_7$. Based on this, the ester was given formula (I) and the lactone formula (II).



(I) (II)

Since more than one (OH) group is present in the free acid molecule, a test for viscinal (OH) groups was made by measuring the exaltation of rotation produced by borax. $(\alpha)_D^{30}$ for free acid = −20°; after saturation with borax = −92°, which is clearly indicative of "the situation of the two hydroxyl groups in the same plane and on the same side of the carbon atoms to which they are bound".[12]

Since it was now evident that the acid from Garcinia is one of the Hydroxycitric acids, (+)-allo -Hydroxycitric lactone was prepared from the calyxes of *Hibiscus Sabdariffa* fruits (according to the procedure recommended by Griebel for purposes of comparison. That the two acids are different is clear from the properties listed in Table 2. The properties of "Garcinia" acid closely resembled those of (-)-hydroxycitric acid described by Martius A sample of our lactone was examined by Dr. C. Marks. It was found that the lactone from Garcinia Cambogia, the lactone of synthetic (-)-Hydroxycitric acid and the mixture of both had the same melting point. Paper chromatography also showed the identity of "Garcinia" acid and synthetic (-)-Hydroxycitric acid.

We have also examined the acid composition of a few related plants. The fruits of *Garcinia atroviridis Griff* and *Garcinia indica Choisy*, which are widely used for culinary purposes in India and Malaya, were found to contain (-)-Hydroxycitric acid as the principal acid. *Hibiscus carmabinus Linn* and *Hibiscus fwcatus Roxb* leaves were found to contain (+)-allo - Hydroxycitric acid as the principal acid. The mildly acidic leaves of these two plants are also used for culinary purposes in India. The fruit calyxes in these were, however, poor in acid content.

Discussion

Martius emphasized the importance of Hydroxycitric acid in plant and animal metabolism at a time when it was still not known to occur in nature. A survey of organic acids in a large number of fruits belonging to the Garcinia genus would doubtless reveal the presence of Hydroxycitric acid in many of them besides those reported here. It is also quite likely that the other two isomers (+)-Hydroxycitric acid and (-)-allo - Hydroxycitric acid would be found in this and other genera. The occurrence of isomers of Hydroxycitric acid in nature may thus be quite wide.

Configuration

The exaltation of optical rotation by borate in "Garcinia" acid is clearly indicative of "vicinal" (OH) groups. The allo-hyroxycitric acid from H. sabakriffa does not show any such increase of rotation. If Hudson's lactone rule is applicable to the Hydroxycitric acids (it is shown to be applicable to the isocitric acids 13), then the absolute configuration of "Garcinia" acid should be (III). Correspondingly, those of the other isomers should be (IV), (V)



(−)-Hydroxycitric acid
Lactone dextro
"*Garcinia acid*"

(III)



(+)-Hydroxycitric acid
Lactone levo

(IV)

[12] J. Bosseren, *Rec. Trav. chim.* 34, 96 (1915).
[13] H. Katsura, *Chem. & Ind. (London)* 1188 (1960).

and (VI). Confirmation of this by optical rotatory dispersion and X-ray diffraction studies would, of course, be necessary.



(−)-allo-Hydroxycitric acid
Lactone levo

(V)



(+)-allo-Hydroxycitric acid
Lactone dextro.
"*Hibiscus acid*"

(VI)

R_M Values

Bate-Smith and Westall[14] have shown that in several series of polyhydroxy compounds, the addition of each (OH) group decreases the R_M value [$(R_M = \log(1/R_f - 1))$] of the compounds by an approximately constant amount. Hulme[15] found this to be true in a related series of acids like succinic, malic and tartaric acids. It is interesting to note that this is not obeyed in the series tricarballylic, citric and hydroxycitric acids (Table 3).

TABLE 3. R_M VALUES OF TRICARBALLYLIC ACID SERIES

Acid	BFW R_M	BFW ΔR_M	PAW R_M	PAW ΔR_M
Tricarballylic	−0·31		0·30	
		0·49		0·13
Citric	0·18		0·63	
		0·25		0·38
Hydroxycitric	0·43		0·25	
		0·57*		0·36*
Allo-hydroxycitric	0·75		0·27	

* Between allo-hydroxycitric and citric acid.

Colour Reaction with Metavandate

Since the R_f of Hydroxycitric acid is very close to tartaric acid in the common solvent systems and it also gives a reddish orange colour with metavandate. It would not be sufficient to

depend on these two data for identification of optically active tartaric acid in plants, as has been done previously.

Biochemistry

The mode of formation of the acid in the plant cell would be a very interesting study. Martius found that only one of the four isomers (-) Hydroxycitric acid could be readily dehydrogenated using cucumber seed or liver homogenate as source of the enzyme. The presence of large quantities of the inactive isomers in plants may then be due to the absence of enzymes which could oxidize them further, as in the case of (+) - tartaric acid. This has to be verified by experimentation. Martius makes the assumption that in the formation of hydroxy glutamic acid, (+) - Hydroxycitric acid has an importance at least corresponding to that of citric or isocitric acids, for the endogenous formation of glutamic acid. It may be possible to prove the biological importance of this substrate, at least in plants.

Acknowledgements—Our grateful thanks are due to Dr. H. B. Vickery of the Connecticut Agricultural Experiment Station for the column chromatographic data, Dr. A. C. Hulme of Ditton Laboratories, Kent, for some of the paper chromatographic data, Dr. B. C. Subba Rao of the National Chemical Laboratories and Dr. D. K. Bannerjee of the Indian Institute of Science for the I.R. data and element analysis. Drs. J. Wolf and C. Griebel of Germany and Hakuji Katsura of Japan have, like the others named before, offered useful suggestions. Our special thanks are due to Dr. C. Martius of Zürich for having confirmed our identification of the acid by comparison with his synthetic acid. We are grateful to Dr. H. A. B. Parpia, Director, and Dr. D. S. Bhatia, Assistant Director, of this Institute, for the keen interest they have shown in this work.

[14] H. B. VICKERY and J. K. PALMER, *J. Biol. Chem.* 207, 275 (1954).

References

E.V. Lippmann Ber. Deut. chem. Ges. 16 (1883), p. 1078.
A. Pawollek Liebigs Ann. Chem. 178 (1875), p. 155.
C. Martius and R. Maue Z. physiol. Chem. 33 (1941), p. 269.
C. Griebel Z. Unters. Lebensm. 77 (1939), p. 561.
C. Griebel Z. Unters. Lebensm. 83 (1942), p. 481.
K.I. Kuriyan and K.C. Pandya J. Indian Chem. Soc. (1931), p. 469.
A. Sreenivasan and R. Venkataraman Curr. Sci. (India) 28 (1959), p. 151.
H.A. Stafford Amer. J. Bot. 46 (1959), p. 347.
Official and Tentative Methods of Analysis, Assn. of Agr. Chem. Washington (1955).
J.K. Palmer Bulletin 589, Connecticut Agri. Exp. Station, New Haven, Conn. (1955).
J.R. Stern, S.P. Colowick and N.O. Kaplan Methods in Enzymol. 3 (1957), p. 428.
J. Boeseken Rec. Trav. chim. 34 (1915), p. 96.
H. Katsura Chem. & Ind. (London) (1960), p. 1188.
E.C. Bate-Smith and R.G. Westall Biochim. Biophys. Acta 4 (1950), p. 427.
A.C. Hulme J. Exp. Botany 2 (1951), p. 298.
H.B. Vickery and J.K. Palmer J. Biol. Chem. 207 (1954), p. 275.

TITLE

THE DIET AND HEALTH BENEFITS OF HCA

(Hydroxycitric Acid)

Dallas Clouatre, Ph.D. and Michael Rosenbaum, MD

Summary

(-)-Hydroxycitric Acid extracted from Garcinia Cambogia is identified as the ingredient which prevents the body from converting carbohydrate calories into fats.
(-)-HCA does this by inhibiting the enzyme ATP citrate lyase, the enzyme which converts citrate into fatty acids and cholesterol in the primary pathway of fat synthesis in the body.

The action of (-)-HCA increases the production and storage of glycogen. It is thought that this stimulates glycoreceptors in the liver, which results in early satiety through signals sent to the brain via the vagus nerve. (-)-HCA is close to citric acid in terms of safety results. (-)-HCA does not alter taste or cause gastric distress or illness. Typical does of (-)HCA in humans for the purposes of losing weight is roughly the equivalent of the rind of half a fruit, which is an insignificant amount considering its common use.

CONCLUSIONS OF ARTICLE

(–)-Hydroxycitric acid is one of the most promising natural products to appear in many years. It has been extensively researched at leading universities and by the pharmaceutical giant Hoffmann-La Roche. Long used in large amounts as a condiment for foods, HCA has no toxicity, either acute or chronic, at any reasonable level of consumption. In this regard it is virtually identical to citric acid, the acid found in oranges, lemons and other citrus fruits. HCA works as a "partitioning agent" with calories obtained from carbohydrates, and, to a lesser extent, those from proteins. It diverts these calories away from the production of fats and toward the production of glycogen.

The health benefits of HCA are many. HCA inhibits lipogenesis, which is the production of fats from excess calories derived from carbohydrates and proteins. As an aspect of its regulation of lipogenesis, HCA lowers the production of both triglycerides and cholesterol. HCA increases the production and storage of glycogen in the liver and other tissues. By doing so, it increases energy levels while giving better control over the appetite. Finally, HCA may increase thermogenesis, the burning of calories for heat.

HCA therefore is potentially of great benefit to dieters and to others who are concerned with controlling their weight, including athletes. It may help individuals with problems with elevated blood fat levels. And in general, because of its effects upon glycogen production and storage, HCA likely will improve energy levels in many individuals who otherwise do not see themselves as having weight or blood fat problems.

HOW THIS STUDY IS RELEVANT TO OUR PRODUCT

The article's conclusion is that (-)HCA is safe, with no toxicity, either acute or chronic based on centuries of consumption by humans and many clinical trials.

The conclusion also goes on to explain the inhibition of lipogenesis, it's appetite suppressant qualities and its possible uses as part of a dietary regime promoting weight loss.

Taken together, these conclusions re-enforce the safety and efficacy of our new dietary ingredient.