

TEAR HERE TO OPEN

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ do not take more than directed

adults and children 12 years and over	take 2 caplets at bedtime or as directed by a doctor
children under 12 years	do not use this adult product in children under 12 years of age; this will provide more than the recommended dose (overdose) and could cause serious health problems

Other information

- do not use if container is open or if tear strip is broken or missing
- store at room temperature
- see container label for lot and expiration date

Inactive ingredients

camauba wax, cellulose, cornstarch, FD&C Blue #1, FD&C Blue #2, hydroxypropyl methylcellulose, magnesium stearate, polyethylene glycol, polysorbate 80, sodium citrate, sodium starch glycolate, titanium dioxide

Questions or comments?

call toll-free 1-877-TYLENOL (1-877-895-3665)

You can feel good knowing **TYLENOL® PM:**

- is non habit-forming
- helps you wake up refreshed (best if taken at least 6 hours before you need to wake up).

The makers of **TYLENOL® PM** do not manufacture store brands.



McNeil Consumer Healthcare
DIVISION OF MCNEIL - PPC, INC.
FORT WASHINGTON, PA 19034 USA
© MCN-PPC, Inc. '01
Visit us at www.tylenol.com or call toll-free
1-877-TYLENOL (1-877-895-3665)

XXXXXXX



Drug Facts

Active ingredient (in each caplet)	Purpose
Acetaminophen 500 mg.....	Pain reliever
Diphenhydramine HCl 25 mg.....	Sleep aid

Uses temporary relief of occasional headaches and minor aches and pains with accompanying sleeplessness

Warnings

Alcohol warning: If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take acetaminophen or other pain relievers/fever reducers. Acetaminophen may cause liver damage.

Do not use

- with any other product containing acetaminophen
- in children under 12 years of age

Ask a doctor before use if you have

- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- glaucoma

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

When using this product

- drowsiness will occur
- avoid alcoholic drinks
- do not drive a motor vehicle or operate machinery

Stop use and ask a doctor if

- sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of serious underlying medical illness.
- new symptoms occur
- redness or swelling is present
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days

Labeling Format Information:

Fonts: Helvetica regular, bold and bold italic.	
Drug Facts: 10 pt	Leading: 6.5 pt
Header: 8 pt	Bullets: 5 pt
Subheader: 6 pt	Barlines: 2.5 pt
Body Text: 6 pt	Hairlines: 0.5 pt
Drug Facts (continued): 8 pt	

Horizontal Scale: 90-100%

Average Kerning: 0