ATTACHMENT 1

To Open, Write Folded on Line, Tear At Slit

Maximum Strength

TYLENOL® SINUS

Pain Reliever
Nasal Decongestant

NON-DROWSY

For Fast Relief of:
Headache Pain
Sinus Pressure
Nasal Congestion

SEE NEW LABEL

2 Geltabs

Active Ingredients: **Purposes:**
(In Each Geltab)
Acetaminophen 500 mg.....Pain Reliever
Pseudoephedrine HCl 30 mg.....Nasal Decongestant
Uses: For the temporary relief of sinus pain and headaches and nasal and sinus congestion.
Do not use if pouch is opened.
Store at room temperature; avoid high humidity and excessive heat 40C (104F).
McNeil Consumer Healthcare, DIVISION OF MCNEIL-PPC, INC.
FORT WASHINGTON, PA 19034 USA © MCN-PPC, Inc. '99

Warnings: Alcohol Warning: If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take acetaminophen or other pain relievers/fever reducers. Acetaminophen may cause liver damage. **Warnings:** Do not take for pain for more than 7 days or for fever for more than 3 days unless directed by a doctor. If pain or fever persists, or gets worse, if new symptoms occur, or if redness or swelling is present, consult a doctor because these could be signs of a serious condition. If nervousness, dizziness or sleeplessness occur, discontinue use and consult a doctor. Do not take this product if you have heart disease, high blood pressure, thyroid disease, diabetes or difficulty in urination due to enlargement of the prostate gland unless directed by a doctor. **Do not exceed recommended dosage.** Keep this and all drugs out of the reach of children. In case of accidental overdose, contact a doctor or poison control center immediately. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. As with any drug, if you are pregnant or nursing a baby, seek the advice of a health professional before using this product. Do not use with other products containing acetaminophen. **Drug Interaction Precaution:** Do not use this product if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you are uncertain whether your prescription drug contains an MAOI, consult a health professional before taking this product. **Directions: Adults and children 12 years of age and older: Take 2 geltabs every 4-6 hours. Do not take more than 8 geltabs in 24 hours, or as directed by a doctor. Children under 12 years: Do not use this adult product in children under 12 years of age. This will provide more than the recommended dose (overdose) and could cause serious health problems. Inactive Ingredients:** Benzyl Alcohol, Butylparaben, Castor Oil, Cellulose, Corn Starch, D&C Yellow 10, Edetate Calcium Disodium, FD&C Blue 1, Gelatin, Hydroxypropyl Methylcellulose, Iron Oxide, Magnesium Stearate, Methylparaben, Propylparaben, Sodium Lauryl Sulfate, Sodium Propionate, Sodium Starch Glycolate, Titanium Dioxide. XXXXXXX

150% OF SIZE