

SPECIMEN LABEL #3 - FOR SHIPMENT OF MEDICATED LAYER FEED FOR APPLICANT'S OWN USE

N-040001 Submitted 12/16/1974

292 Approved 5/31/1973

FOR USE ON: BULK FEED PLACARD

Blue Bird Layer Ration (50 lb bag) own use FEED BAG TAG

Type C Med. Feed

50 LBS. NET WEIGHT

BLUE BIRD LAYER RATION

MEDICATED

FOR THE CONTROL OF CAPILLARY WORM (*Capillaria obsignata*)

AND AS AN AID IN THE CONTROL OF COMMON ROUNDWORM (*Ascaridia galli*)

AND CECAL WORM (*Heterakis gallinae*) IN LAYING FLOCKS

ACTIVE DRUG INGREDIENT:

coumaphos 0.003%

DIRECTIONS FOR USE

A diagnosis by competent personnel is essential. Administer continuously as the total daily feed ration for 14 days. When reinfection occurs, treatment may be repeated but not sooner than three weeks after the end of the previous treatment. Conditions of the flock and production records should be carefully evaluated before treatment with this medicated ration. As a general rule, all medications should be avoided while birds are near peak production, since interruption of normal feeding practices may upset the flock and lower egg production.

WARNINGS AND LIMITATIONS

Should be the sole source of any medication. Do not feed to chickens within 10 days of vaccination or other conditions of stress. Treatment of colored breeds of commercial layers should be avoided while in production since these breeds appear to be more sensitive to coumaphos than white breeds.

WARNING: Coumaphos is a cholinesterase inhibitor. Do not use this product in animals simultaneously or within a few days before or after treatment with or exposure to cholinesterase inhibiting drugs, pesticides or chemicals.

Symptoms of overdose are difficult to identify because numerous other conditions cause similar symptoms. In case of abnormal behavior of the chickens, discontinue feeding medicated feed and consult your veterinarian or diagnostic laboratory.

FOR OWN USE - NOT FOR SALE

Manufactured by

BLUE BIRD FEED MILLS

BLUE BIRD, MARYLAND 21043