

**THIS BLUEBIRD LABEL IS INTENDED FOR USE ONLY WITH
Zilpaterol, Monensin, and Tylosin Finishing Cattle Feed
Type C Medicated Complete Feed**

**HEIFER TYPE C MEDICATED FEED – BAG or BULK
For Beef and Dairy Heifers**

Caution: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

Heifers Fed in Confinement for Slaughter: For increased rate of weight gain, improved feed efficiency, and suppression of estrus (heat).

ACTIVE DRUG INGREDIENT

Melengestrol acetate..... (0.0000276-0.00022%)*
Each pound contains (0.125-1.0) mg melengestrol acetate..... (select one)*

GUARANTEED ANALYSIS

Crude Protein, not less than	_____	%
Non-Protein Nitrogen (NPN) ¹ , not more than.....	_____	%
Crude Fat, not less than	_____	%
Crude Fiber, not more than	_____	%
Calcium, not less than.....	_____	%
Calcium, not more than	_____	%
Phosphorus, not less than.....	_____	%
Salt ² , not less than	_____	%
Salt ² , not more than	_____	%
Sodium ³ , not less than	_____	%
Sodium ³ , not more than	_____	%
Potassium, not less than.....	_____	%
Vitamin A ^{2,4} , not less than	_____	I.U./lb

¹When added

²If added.

³Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

⁴Other than precursors of Vitamin A.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

*Final printed label of Type C Medicated Feed must bear a single drug concentration of this drug.

FEEDING DIRECTIONS

Must be top dressed or mixed with a Type C feed containing 6.8 g/ton zippaterol, 100 to 40 g/ton monensin, and 8-10 g/ton tylosin.

Feed at the rate of 0.5 to 2.0 lb/head/day (specify one level) to provide 0.25 to 0.5 mg melengestrol acetate/head/day (specify one level). Feed melengestrol acetate in this combination for the final 20 to 40 days on feed.

CAUTION: Melengestrol acetate is not effective in steers and spayed heifers. Withdrawal periods of three to five days or more should be avoided to prevent the possibility that the heifers may come into estrus (heat) at loading time.

Do not use in any finished feed (supplement, concentrate or complete feed) containing in excess of 2% bentonite.

MANUFACTURED BY
BLUE BIRD FEED MILL
Any town, USA 12345

Net Weight lb (kg) on bag or bulk

Batch (Lot or Control) No.:

ANADA 200-544 May 2016