

Label 1: Replacement Chickens Type C Medicated Feed Coccidiostat	
Indications for Use	
For the development of active immunity to coccidiosis.	
Active Drug Ingredient	
Amprolium.....	0.004-0.0125%
Guaranteed Analysis	
Crude Protein, not less than	xx.x%
Crude Fat, not less than	xx.x%
Crude Fiber, not more than	xx.x%
Lysine, not less than	xx.x%
Methionine, not less than	xx.x%
Calcium, min and max	xx.x%
Phosphorus, not less than	xx.x%
Salt, min and max (if added)	xx.x%
Sodium, min and max (when total Sodium exceed the furnished by the maximum salt guarantee)	xx.x%
Ingredients	
Ingredients as defined by AAFCO	
Feeding Directions	
Feed continuously until onset of production.	
Caution	
Use as the sole source of amprolium.	
Do not change the litter while giving this feed unless absolutely necessary.	
If losses exceed 0.5% in a 2-day period, obtain an accurate diagnosis, and follow the instructions of your veterinarian or poultry pathologist.	
Losses may result from intercurrent disease or other conditions affecting drug intake which can contribute to the virulence of coccidiosis under field conditions.	
In replacement flocks the grower must expect that excessive exposure to one or more species may overwhelm the drug in some flocks and prompt treatment will be required.	
Do not use in feeds containing bentonite.	

Warning

Keep this and all drugs out of the reach of children.

Manufactured By

Blue Bird Feed Mill
Robin, Indiana 00000

Net Weight __ lbs (__ kg)
Bag or Bulk