

**Acanya**  
(Clindamycin Phosphate and Benzoyl Peroxide) Gel, 1.2%/2.5%  
**Ready-to-Use 50g Pump**

[Home](#) | [Contact us](#) | [Full prescribing information](#) | [Search](#)

*Power to please*

[About Acanya Gel](#) | [About acne](#) | [Before-and-after](#) | [Special offers](#) | [Healthcare professionals](#)



**Acanya® Gel**—the once-daily topical prescription treatment with the power to treat your acne without overpowering your skin.

**Acanya Gel is proven to work**

Studies showed that Acanya Gel was proven to treat blackheads, whiteheads, and inflamed acne on the face.

**Acanya Gel is gentle to skin, too**

The water-based gel contains no alcohol or detergents that may contribute to stinging or burning. Fragrance-free Acanya Gel goes on smoothly, disappears fast, and leaves no residue.

Now in a convenient pump.



*Get Instant Savings*

Patients pay no more than \$25 every time they fill a prescription for Acanya Gel.

*Tell a friend about Acanya Gel* >  
*and receive a coupon for \$1 off your next purchase of CeraVe® Facial Moisturizer!*

[View instructions for using Acanya Gel](#)



**Indication and Important Safety Information:** Acanya® Gel is indicated for the topical treatment of acne vulgaris in patients 12 years of age or older. Side effects may include redness, scaling, itching, burning, and stinging. Do not use Acanya Gel if you are allergic to clindamycin, or if you have a history of inflammation of the colon (colitis). Clindamycin, the antibiotic in Acanya Gel, may cause diarrhea. If you experience diarrhea, stop using Acanya Gel immediately and contact your doctor. Avoid using tanning beds or sun lamps, and limit your time in sunlight after applying Acanya Gel. Avoid applying Acanya Gel in mouth, eyes, or nose, or on lips.

[Please see full prescribing information for Acanya Gel.](#) >

**VALEANT** © 2011 CORIA Laboratories, a division of Valeant Pharmaceuticals North America LLC

[Terms of use](#) | [Privacy policy](#) | [Site map](#)

**Acanya**  
(Clindamycin Phosphate and  
Benzoyl Peroxide) Gel, 1.2%/2.5%  
Ready-to-Use 50g Pump

[Home](#) | [Contact us](#) | [Full prescribing information](#) | [Search](#)

*Power to please*

[About Acanya Gel](#) | [About acne](#) | [Before-and-after](#) | [Special offers](#) | [Healthcare professionals](#)



#### Talk to your doctor about the right acne medication for you

The more you know about your acne, the better your doctor will be able to help you with treatment. Below are questions to help with your discussion. Before using Acanya Gel, be sure to read the patient information and application instructions. Tell your doctor immediately about any side effect that bothers you or that does not go away.

##### What type of acne do I have?

Acanya Gel is approved to treat mild, moderate, and severe acne. This can include:  
—Noninflammatory acne (blackheads and whiteheads)  
—Inflammatory acne (pimples and pustules)

##### How often will I need to use my acne medication?

Acanya Gel is applied once a day.

##### How long does it take before I see results?

In clinical studies, some patients using Acanya Gel daily saw improvement as soon as 2 weeks, which continued throughout a 12-week treatment period. Both the patients and their doctors continued to see improvement in the patients' acne during that time.

##### Will my acne medication irritate my skin?

Acanya Gel is water based, with no alcohol or detergents that may sting or burn your skin. In clinical studies, infrequent side effects seen were itching, burning, stinging, redness, or flaking skin. However, no patients stopped using Acanya Gel due to these side effects.

##### Can I wear makeup or sunscreen with my acne medication?

Skin treated with Acanya Gel should be protected from sunlight. Ask your doctor about the proper use of sunscreens, moisturizers, and cosmetics with Acanya Gel.

**Indication and Important Safety Information:** Acanya® Gel is indicated for the topical treatment of acne vulgaris in patients 12 years of age or older. Side effects may include redness, scaling, itching, burning, and stinging. Do not use Acanya Gel if you are allergic to clindamycin, or if you have a history of inflammation of the colon (colitis). Clindamycin, the antibiotic in Acanya Gel, may cause diarrhea. If you experience diarrhea, stop using Acanya Gel immediately and contact your doctor. Avoid using tanning beds or sun lamps, and limit your time in sunlight after applying Acanya Gel. Avoid applying Acanya Gel in mouth, eyes, or nose, or on lips.

[Please see full prescribing information for Acanya Gel.](#) >

© 2011 CORIA Laboratories, a division of Valeant Pharmaceuticals North America LLC

[Terms of use](#) | [Privacy policy](#) | [Site map](#)

*Get Instant Savings*

Patients pay no more than \$25 every time they fill a prescription for Acanya Gel.



*Click here*  
to see before-and-after photos >



#### View instructions for using Acanya Gel



#### Download brochure for Acanya Gel



**Acanya**  
(Clindamycin Phosphate and  
Benzoyl Peroxide) Gel, 1.2%/2.5%  
**Ready-to-Use 50g Pump**

[Home](#) | [Contact us](#) | [Full prescribing information](#) | [Search](#)

*Power to please*

[About Acanya Gel](#) | [About acne](#) | [Before-and-after](#) | [Special offers](#) | [Healthcare professionals](#)



The following links can help you learn more about acne and caring for your skin.

**The American Academy of Dermatology (AAD) Acne Overview**

A brief overview of acne and medical treatment options provided to patients by the AAD, an organization of medical professionals.

**Medline Acne Tutorial**

This easy-to-use online tutorial from the National Institutes of Health provides in-depth information about acne symptoms, treatment, and tips for skin care.

[En Español](#)

**Q&A About Acne From NIAMS**

A medical perspective on acne for patient audiences, provided by the National Institute of Arthritis and Musculoskeletal and Skin Diseases.

**Mayo Clinic: Acne**

This overview from the nationally renowned Mayo Clinic covers acne causes and treatments—including adult acne care and prevention.

**Acne and Women's Health**

Information about acne triggers and skin care specific to female patients, provided by the US Department of Health and Human Services.

*Get Instant Savings*

Patients pay no more than \$25 every time they fill a prescription for Acanya Gel.



*Click here*

to see before-and-after photos >



**View instructions for using Acanya Gel**



**Indication and Important Safety Information:** Acanya<sup>®</sup> Gel is indicated for the topical treatment of acne vulgaris in patients 12 years of age or older. Side effects may include redness, scaling, itching, burning, and stinging. Do not use Acanya Gel if you are allergic to clindamycin, or if you have a history of inflammation of the colon (colitis). Clindamycin, the antibiotic in Acanya Gel, may cause diarrhea. If you experience diarrhea, stop using Acanya Gel immediately and contact your doctor. Avoid using tanning beds or sun lamps, and limit your time in sunlight after applying Acanya Gel. Avoid applying Acanya Gel in mouth, eyes, or nose, or on lips.

Please see full prescribing information for Acanya Gel. >

© 2011 CORIA Laboratories, a division of Valeant Pharmaceuticals North America LLC

[Terms of use](#) | [Privacy policy](#) | [Site map](#)

**Acanya**  
(Clindamycin Phosphate and  
Benzoyl Peroxide) Gel, 1.2%/2.5%  
**Ready-to-Use 50g Pump**

[Home](#) | [Contact us](#) | [Full prescribing information](#) [Search](#)



*Power to please*

[About Acanya Gel](#) [About acne](#) [Before-and-after](#) [Special offers](#)  [Healthcare professionals](#)



**Power to treat**

As demonstrated in these patient photos following 12 weeks of once-daily treatment with Acanya® Gel alone:



These unretouched photos represent actual clinical trial experience. As with all treatments, results may vary from person to person.

*Get Instant Savings*

Patients pay no more than \$25 every time they fill a prescription for Acanya Gel.

[View instructions for using Acanya Gel](#)



**Indication and Important Safety Information:** Acanya® Gel is indicated for the topical treatment of acne vulgaris in patients 12 years of age or older. Side effects may include redness, scaling, itching, burning, and stinging. Do not use Acanya Gel if you are allergic to clindamycin, or if you have a history of inflammation of the colon (colitis). Clindamycin, the antibiotic in Acanya Gel, may cause diarrhea. If you experience diarrhea, stop using Acanya Gel immediately and contact your doctor. Avoid using tanning beds or sun lamps, and limit your time in sunlight after applying Acanya Gel. Avoid applying Acanya Gel in mouth, eyes, or nose, or on lips.

Please see full prescribing information for Acanya Gel. >



**Acanya**  
(Clindamycin Phosphate and Benzoyl Peroxide) Gel, 1.2%/2.5%  
Ready-to-Use 50g Pump

[Home](#) | [Contact us](#) | [Full prescribing information](#) | [Search](#)



*Power to please*

[About Acanya Gel](#) | [About acne](#) | [Before-and-after](#) | [Special offers](#) | [Healthcare professionals](#)



Save money on your Acanya® Gel prescription



**Unlimited Use!**

**If you have been prescribed Acanya Gel:**  
Eligible patients\* pay no more than \$25 for each Acanya Gel prescription.

**Download your Acanya Gel Instant Savings Card here.**

\*Offer is good only for US residents and is subject to the terms and eligibility requirements stated on the card.

[Click here](#)  
to see before-  
and-after photos >



[View instructions for using Acanya Gel](#)



**Indication and Important Safety Information for Acanya Gel:** Acanya® Gel is indicated for the topical treatment of acne vulgaris in patients 12 years of age or older. Side effects may include redness, scaling, itching, burning, and stinging. Do not use Acanya Gel if you are allergic to clindamycin, or if you have a history of inflammation of the colon (colitis). Clindamycin, the antibiotic in Acanya Gel, may cause diarrhea. If you experience diarrhea, stop using Acanya Gel immediately and contact your doctor. Avoid using tanning beds or sun lamps, and limit your time in sunlight after applying Acanya Gel. Avoid applying Acanya Gel in mouth, eyes, or nose, or on lips.

[Please see full prescribing information for Acanya Gel.](#) >

© 2011 CORIA Laboratories, a division of Valeant Pharmaceuticals North America LLC

[Terms of use](#) | [Privacy policy](#) | [Site map](#)